Package 'FANTOM3and4CAGE'

April 8, 2015

Title CAGE data from FANTOM3 and FANTOM4 projects
Version 1.1.1
Date 02-06-2013
Author Vanja Haberle, Department of Biology, University of Bergen, Norway <vanja.haberle@bio.uib.no></vanja.haberle@bio.uib.no>
Maintainer Vanja Haberle <vanja.haberle@bio.uib.no></vanja.haberle@bio.uib.no>
Depends R (>= $2.15.0$)
Description CAGE (Cap Analysis Gene Expression) data from FANTOM3 and FANTOM4 projects produced by RIKEN Omics Science Center.
License GPL-3
biocViews ExperimentData, TimeCourse, Transcription, HighThroughputSequencing
R topics documented:
FANTOMhumanSamples FANTOMmouseSamples FANTOMtimecourseCAGEhuman FANTOMtimecourseCAGEmouse FANTOMtissueCAGEhuman FANTOMtissueCAGEmouse FANTOMtissueCAGEmouse
Index
FANTOMhumanSamples List of FANTOM CAGE samples for human

Description

A data.frame with information on all human CAGE samples from FANTOM3 and FANTOM4 projects available in this package.

Usage

```
data(FANTOMhumanSamples)
```

Details

FANTOMhumanSamples is a data.frame with 3 columns:

dataset: the name of the dataset that can be loaded using data() function group: the name of the group of samples that originate from the same tissue (*e.g.* blood) sample: the name of the specific sample

Source

The annotation of CAGE samples was obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860.

Examples

```
data(FANTOMhumanSamples)
head(FANTOMhumanSamples)
```

FANTOMmouseSamples

List of FANTOM CAGE samples for mouse

Description

A data.frame with information on all mouse CAGE samples from FANTOM3 and FANTOM4 projects available in this package.

Usage

```
data(FANTOMmouseSamples)
```

Details

FANTOMmouseSamples is a data. frame with 3 columns:

dataset: the name of the dataset that can be loaded using data() function group: the name of the group of samples that originate from the same tissue (*e.g.* brain) sample: the name of the specific sample

Source

The annotation of CAGE samples was obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860.

Examples

data(FANTOMmouseSamples)
head(FANTOMmouseSamples)

FANTOMtimecourseCAGEhuman

Human timecourse CAGE data produced in FANTOM4 project

Description

This dataset contains all transcription start sites (TSSs) detected by CAGE and their frequency of usage in the human THP-1 cells induction timecourse from FANTOM4 project.

Usage

data(FANTOMtimecourseCAGEhuman)

Details

FANTOMtimecourseCAGEhuman is a list with only one element (timecourse) named THP-1_monocytic_induction. This element is a data. frame with genomic positions of TSSs detected by CAGE and frequency of their usage in various timepoints during stimulation of human THP-1 myelomonocytic leukemia cells by phorbol myristate acetate (PMA). First three columns contain the chromosome (chr), the 1-based coordinate (pos) and the strand (strand) of the TSS, and the following columns contain the number of CAGE tags supporting that TSS in each specified timepoint.

Source

The CAGE detected TSSs (mapped to the hg18 human genome assembly) and numbers of CAGE tags for THP-1 monocytic induction were obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/) The data were originally published by The FANTOM Consortium (Suzuki *et al.*, Nature Genetics 2009).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860. Suzuki *et al.* (2009) The transcriptional network that controls growth arrest and differentiation in a human myeloid leukemia cell line, *Nature Genetics* **41**:553-562.

Examples

```
data(FANTOMtimecourseCAGEhuman)
names(FANTOMtimecourseCAGEhuman)
head(FANTOMtimecourseCAGEhuman[["THP-1_monocytic_induction"]])
```

FANTOMtimecourseCAGEmouse

Mouse timecourse CAGE data produced in FANTOM projects

Description

This dataset contains all transcription start sites (TSSs) detected by CAGE and their frequency of usage in two mouse timecourses: adipogenic induction of DFAT-D1 preadipocytes and liver under constant darkness condition.

Usage

data(FANTOMtimecourseCAGEmouse)

Details

FANTOMtimecourseCAGEmouse is a list with two elements (timecourses) named adipogenic_induction and liver_under_constant_darkness. Each element is a data. frame with genomic positions of TSSs detected by CAGE and frequency of their usage in various timepoints during timecourse. First three columns contain the chromosome (chr), the 1-based coordinate (pos) and the strand (strand) of the TSS, and the following columns contain the number of CAGE tags supporting that TSS in each specified timepoint.

Source

The CAGE detected TSSs (mapped to the mm9 mouse genome assembly) and numbers of CAGE tags for both timecourses were obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/). The data were originally published by The FANTOM Consortium (Carninci *et al.*, Science 2005).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860.

Carninci *et al.* (2005) The Transcriptional Landscape of the Mammalian Genome, *Science* **309**(5740):1559-1563.

Examples

```
data(FANTOMtimecourseCAGEmouse)
names(FANTOMtimecourseCAGEmouse)
head(FANTOMtimecourseCAGEmouse[["adipogenic_induction"]])
```

FANTOMtissueCAGEhuman CAGE data produced in FANTOM projects for various human tissues and cell lines

Description

This dataset contains all transcription start sites (TSSs) detected by CAGE and their frequency of usage in the various human tissues and cell lines generated in FANTOM3 and FANTOM4 projects.

Usage

data(FANTOMtissueCAGEhuman)

Details

FANTOMtissueCAGEhuman is a list of groups of human samples, which are grouped by the tissue of origin. Each element is named according to the corresponding tissue and it is a data.frame with genomic positions of TSSs detected by CAGE and frequency of their usage in various samples originating from that tissue. First three columns in every data.frame contain the chromosome (chr), the 1-based coordinate (pos) and the strand (strand) of the TSS, and the following columns contain the number of CAGE tags supporting that TSS in each specific sample. These columns are named according to the corresponding sample.

Source

The CAGE detected TSSs (mapped to the hg18 human genome assembly) and numbers of CAGE tags were obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/). The data were originally published by The FANTOM Consortium (Carninci *et al.*, Science 2005, and Carninci *et al.*, Nature Genetics 2006, and Faulkner *et al.*, Nature Genetics 2009).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860.

Carninci *et al.* (2005) The transcriptional landscape of the mammalian genome, *Science* **309**:1559-1563.

Carninci *et al.* (2006) Genome-wide analysis of mammalian promoter architecture and evolution, *Nature Genetics* **6**:626-635.

Faulkner et al. (2009) The regulated retrotransposon transcriptome of mammalian cells, *Nature Genetics* **41**:563-571.

Examples

data(FANTOMtissueCAGEhuman)
names(FANTOMtissueCAGEhuman)
head(FANTOMtissueCAGEhuman[["liver"]])

FANTOMtissueCAGEmouse CAGE data produced in FANTOM projects for various mouse tissues and cell lines

Description

This dataset contains all transcription start sites (TSSs) detected by CAGE and their frequency of usage in the various mouse tissues and cell lines generated in FANTOM3 and FANTOM4 projects.

Usage

data(FANTOMtissueCAGEmouse)

Details

FANTOMtissueCAGEmouse is a list of groups of mouse samples, which are grouped by the tissue of origin. Each element is named according to the corresponding tissue and it is a data.frame with genomic positions of TSSs detected by CAGE and frequency of their usage in various samples originating from that tissue. First three columns in every data.frame contain the chromosome (chr), the 1-based coordinate (pos) and the strand (strand) of the TSS, and the following columns contain the number of CAGE tags supporting that TSS in each specific sample. These columns are named according to the corresponding sample.

Source

The CAGE detected TSSs (mapped to the mm9 mouse genome assembly) and numbers of CAGE tags were obtained from the FANTOM web resource (http://fantom.gsc.riken.jp/4/download/). The data were originally published by The FANTOM Consortium (Carninci *et al.*, Science 2005, and Faulkner *et al.*, Nature Genetics 2009).

References

Kawaji *et al.* (2010) Update of the FANTOM web resource: from mammalian transcriptional land-scape to its dynamic regulation, *Nucleic Acids Research* **39**:D856-D860.

Carninci et al. (2005) The transcriptional landscape of the mammalian genome, Science 309:1559-1563

Faulkner et al. (2009) The regulated retrotransposon transcriptome of mammalian cells, *Nature Genetics* **41**:563-571.

Examples

data(FANTOMtissueCAGEmouse)
names(FANTOMtissueCAGEmouse)
head(FANTOMtissueCAGEmouse[["lung"]])

Index

*Topic datasets FANTOMhumanSamples, 1 FANTOMmouseSamples, 2 FANTOMtimecourseCAGEhuman, 3 FANTOMtimecourseCAGEmouse, 4 FANTOMtissueCAGEhuman, 5 FANTOMtissueCAGEmouse, 6 FANTOMhumanSamples, 1 FANTOMmouseSamples, 2 FANTOMtimecourseCAGEhuman, 3 FANTOMtimecourseCAGEhuman, 3 FANTOMtissueCAGEhuman, 5 FANTOMtissueCAGEhuman, 5 FANTOMtissueCAGEmouse, 6