# Package 'curatedCRCData'

# April 16, 2024

| April 10, 2024   |
|--|
| Type Package   |
| Title Colorectal Cancer Gene Expression Analysis   |
| Version 2.34.0   |
| <b>Date</b> 2015-07-09   |
| Author Princy Parsana, Markus Riester, Curtis Huttenhower, Levi Waldron  |
| Maintainer Princy Parsana <princyparsana@jhu.edu></princyparsana@jhu.edu>  |
| <b>Description</b> The curatedCRC package provides relevant functions and data for gene expression analy sis in patients with colorectal cancer. |
| <b>Depends</b> R (>= $2.10.0$ ), nlme  |
| Imports BiocGenerics   |
| <b>Suggests</b> survival, RUnit, metafor, genefilter, logging, sva, xtable, futile.logger, BiocStyle   |
| Requires affy  |
| License Artistic-2.0   |
| Namespace auto   |
| biocViews Colorectal, Cancer, TCGA, ExperimentData, RNAExpressionData  |
| <pre>URL https://bitbucket.org/biobakery/curatedcrcdata</pre>  |
| git_url https://git.bioconductor.org/packages/curatedCRCData   |
| git_branch RELEASE_3_18  |
| git_last_commit 3df2048  |
| git_last_commit_date 2023-10-24  |
| Repository Bioconductor 3.18   |
| Date/Publication 2024-04-16  |
| R topics documented:   |
| curatedCRCData-package   |

|       | GSE12945_eset           |
|-------|-------------------------|
|       | GSE13067_eset           |
|       | GSE13294_eset           |
|       | GSE14095_eset           |
|       | GSE14333_eset           |
|       | GSE16125.GPL5175_eset   |
|       | GSE17536_eset           |
|       | GSE17537_eset           |
|       | GSE17538.GPL570_eset    |
|       | GSE18105_eset           |
|       | GSE2109_eset            |
|       | GSE21510_eset           |
|       | GSE21815_eset           |
|       | GSE24549.GPL5175_eset   |
|       | GSE24550.GPL5175_eset   |
|       | GSE2630_eset            |
|       | GSE26682.GPL570_eset    |
|       | GSE26682.GPL96_eset     |
|       | GSE26906_eset           |
|       | GSE27544_eset           |
|       | GSE28702_eset           |
|       | GSE3294_eset            |
|       | GSE33113_eset           |
|       | GSE39582_eset           |
|       | GSE3964_eset            |
|       | GSE4045_eset            |
|       | GSE4526_eset            |
|       | GSE45270_eset           |
|       | TCGA.COAD_eset          |
|       | TCGA.READ_eset          |
|       | TCGA.RNASeqV2.READ_eset |
|       | TCGA.RNASeqV2_eset      |
|       |                         |
| Index | 105                     |

curatedCRCData-package

Clinically Annotated Data for the CRC Cancer Transcriptome

# Description

The curatedCRCData package provides manually curated clinical data, uniformly processed expression data, and convenience functions for gene expression analysis in patients with colorectal cancer.

Note: For staging, the "summary stage" was curated as using information on the pathologic T, N, and M stages, such that "early" = T stage of T is through T3 and N0 and M0 and "late" = T stage of 4 and/or any N>0 or M>0 GSE11237\_eset 3

#### **Details**

Package: curatedCRCData

Type: Package
Version: 2.34.0
Date: 2015-07-09
License: Artistic-2.0
Depends: R (>= 2.10.0), affy

#### Author(s)

Princy Parsana, Markus Riester, Levi Waldron

Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute, Harvard School of Public Health

Maintainer: Princy Parsana <pri>princyparsana@jhu.edu>

# **Examples**

```
##List all datasets:
data(package="curatedCRCData")
##
```

GSE11237\_eset

Celecoxib pre-treatment in human colorectal adenocarcinoma patients is associated with gene expression alterations suggestive of diminished cellular proliferation.

#### **Description**

Cancer cells treated with the cyclooxygenase-2 inhibitor celecoxib show growth inhibition and induced apoptosis. This study was conducted to determine if the same processes are relevant to celecoxib's effects on human colorectal adenocarcinomas treated in vivo. A cohort of 23 patients with primary colorectal adenocarcinomas was randomised to receive a 7-d course of celecoxib (400mg b.i.d.) or no drug prior to surgical resection. Gene expression profiling was performed on resected adenocarcinomas from the cohort of patients. Using fold change (>1.5) and p-value (<0.05) cutoffs, 190 genes were differentially expressed between adenocarcinomas from patients receiving celecoxib and those that did not. The celecoxib pre-treated samples showed decreased expression levels in multiple genes involved in cellular lipid and glutathione metabolism; changes associated with diminished cellular proliferation. Celecoxib pre-treatment for 7 d in vivo is associated with alterations in colorectal adenocarcinoma gene expression which are suggestive of diminished cellular proliferation.

4 GSE11237\_eset

# Usage

```
data( GSE11237_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Auman JT, Church R, Lee SY, Watson MA, Fleshman JW, Mcleod HL.??Celecoxib pre-treati
 Laboratory: Auman, Mcleod 2008
 Contact information:
 Title: Celecoxib pre-treatment in human colorectal adenocarcinoma patients is associated with gene exp
 PMIDs: 18653328
  Abstract: A 147 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      Affymetrix Human Genome U95 Version 2 Array
   platform_shorttitle:
      Affymetrix HG_U95Av2
   platform_summary:
      hgu95av2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL8300
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: rma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: AADAC AAK1 ... ZZZ3 (8933 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

# **Details**

GSE11237\_eset 5

```
-----
alt_sample_name:
  Length Class Mode
     23 character character
sample_type:
tumor
  23
summarygrade:
high low
 4 19
G:
1 2 3
3 16 4
summarystage:
early late
 12 11
T:
1 2 3 4
1 3 16 3
N:
0 1 2
15 7 1
M:
0 1
17 6
location:
  Length
           Class
                    Mode
     23 character character
summarylocation:
1 r
9 14
gender:
f m
13 10
stageall:
```

1 2 3 4

```
4 9 4 6

batch:

2003-08-19 2003-08-28

15 8

preop_drug_treatment:
n y
12 11

preop_drug_name:
celecoxib NA's
11 12

uncurated_author_metadata:
   Length Class Mode
   23 character character
```

GSE12225.GPL3676\_eset Integrating chromosomal aberrations and gene expression profiles to dissect rectal tumorigenesis.

#### **Description**

Accurate staging of rectal tumors is essential for making the correct treatment choice. In a previous study, we found that loss of 17p, 18q and gain of 8q, 13q and 20q could distinguish adenoma from carcinoma tissue and that gain of 1q was related to lymph node metastasis. In order to find markers for tumor staging, we searched for candidate genes on these specific chromosomes. We performed gene expression microarray analysis on 79 rectal tumors and integrated these data with genomic data from the same sample series. We performed supervised analysis to find candidate genes on affected chromosomes and validated the results with qRT-PCR and immunohistochemistry.Integration of gene expression and chromosomal instability data revealed similarity between these two data types. Supervised analysis identified up-regulation of EFNA1 in cases with 1q gain, and EFNA1 expression was correlated with the expression of a target gene (VEGF). The BOP1 gene, involved in ribosome biogenesis and related to chromosomal instability, was over-expressed in cases with 8q gain. SMAD2 was the most down-regulated gene on 18q, and on 20q, STMN3 and TGIF2 were highly up-regulated. Immunohistochemistry for SMAD4 correlated with SMAD2 gene expression and 18q loss. On basis of integrative analysis this study identified one well known CRC gene (SMAD2) and several other genes (EFNA1, BOP1, TGIF2 and STMN3) that possibly could be used for rectal cancer characterization.

#### Usage

```
data( GSE12225.GPL3676_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Lips EH, van Eijk R, de Graaf EJ, Oosting J et al.??Integrating chromosomal aberrati
 Laboratory: Lips, Morreau 2008
 Contact information:
 Title: Integrating chromosomal aberrations and gene expression profiles to dissect rectal tumorigeness
 URL:
 PMIDs: 18959792
 Abstract: A 221 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      NKI-CMF Homo sapiens 35k oligo array
   platform_shorttitle:
      NA
   platform_summary:
      nki-cmf homo sapiens 35k oligo array
   platform_manufacturer:
      Central Microarray Facility, NKI Amsterdam
   platform_distribution:
      non-commercial
   platform_accession:
      GPL3676
   platform_technology:
      spotted oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 15E1.2///TRIAP1///SFRS9///DYNLL1///COX6A1
    1810006K21Rik///C11orf10 ... ZZZ3 (17015 total)
 varLabels: probeset gene
  varMetadata: labelDescription
```

#### **Details**

```
Length
             Class
                         Mode
       42 character character
sample_type:
tumor
   42
primarysite:
re
42
summarystage:
early late
   30 12
T:
1 2 3
14 27 1
 0 1 2
30 6 6
M:
 0
42
location:
rectum
    42
\verb|summarylocation|:
1
42
stageall:
1 3
29 13
{\tt preop\_drug\_treatment:}
n
42
uncurated_author_metadata:
            Class
   Length
       42 character character
```

GSE12945\_eset 9

GSE12945\_eset An expression module of WIPF1-coexpressed genes identifies patients with favorable prognosis in three tumor types.

#### Description

Wiskott-Aldrich syndrome (WAS) predisposes patients to leukemia and lymphoma. WAS is caused by mutations in the protein WASP which impair its interaction with the WIPF1 protein. Here, we aim to identify a module of WIPF1-coexpressed genes and to assess its use as a prognostic signature for colorectal cancer, glioma, and breast cancer patients. Two public colorectal cancer microarray data sets were used for discovery and validation of the WIPF1 co-expression module. Based on expression of the WIPF1 signature, we classified more than 400 additional tumors with microarray data from our own experiments or from publicly available data sets according to their WIPF1 signature expression. This allowed us to separate patient populations for colorectal cancers, breast cancers, and gliomas for which clinical characteristics like survival times and times to relapse were analyzed. Groups of colorectal cancer, breast cancer, and glioma patients with low expression of the WIPF1 co-expression module generally had a favorable prognosis. In addition, the majority of WIPF1 signature genes are individually correlated with disease outcome in different studies. Literature gene network analysis revealed that among WIPF1 co-expressed genes known direct transcriptional targets of c-myc, ESR1 and p53 are enriched. The mean expression profile of WIPF1 signature genes is correlated with the profile of a proliferation signature. The WIPF1 signature is the first microarray-based prognostic expression signature primarily developed for colorectal cancer that is instrumental in other tumor types: low expression of the WIPF1 module is associated with better prognosis.

# Usage

```
data( GSE12945_eset )
```

```
experimentData(eset):
Experiment data

Experimenter name: Staub E, Groene J, Heinze M, Mennerich D et al.??An expression module of WIPF1-coexp Laboratory: Staub, Rosenthal 2009
Contact information:
Title: An expression module of WIPF1-coexpressed genes identifies patients with favorable prognosis in URL:
PMIDs: 19399471

Abstract: A 241 word abstract is available. Use 'abstract' method.
Information is available on: preprocessing notes:
   platform_title:
      [HG-U133A] Affymetrix Human Genome U133A Array
   platform_shorttitle:
      Affymetrix HG-U133A
```

10 GSE12945\_eset

```
platform_summary:
         hgu133a
      platform_manufacturer:
         Affymetrix
      platform_distribution:
         commercial
      platform_accession:
         GPL96
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1CF A2M ... ZZZ3 (12986 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 12986 features, 62 samples
   Platform type: hgu133a
   Overall survival time-to-event summary (in years):
   Call: survfit(formula = Surv(time, cens) ~ -1)
             n.max n.start events median 0.95LCL 0.95UCL
   records
        62
               62
                      62
                           12
                                      NA
                                              NA
    _____
   Available sample meta-data:
   -----
   alt_sample_name:
      Length
                Class
                           Mode
          62 character character
   sample_type:
   tumor
      62
   primarysite:
   co re
   29 33
   summarygrade:
   high low
```

GSE12945\_eset 11

```
G:
2 3
31 31
summarystage:
early late
  36 26
T:
2 3 4
16 42 4
N:
0 1 2
36 14 12
M:
0 1 X
56 5 1
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median Mean 3rd Qu.
                                      Max.
  38.00 59.00 65.00 64.45 73.75
                                     87.00
recurrence_status:
norecurrence recurrence
                              NA's
                              7
              8
         47
days_to_death:
  Min. 1st Qu. Median
                        Mean 3rd Qu.
                                      Max.
   210 1005 1395
                      1267 1620
                                      1920
vital_status:
deceased living
     12
location:
  Length
            Class
                      Mode
     62 character character
summarylocation:
  1 r NA's
 48 13 1
gender:
f m
```

31 31

12 GSE13067\_eset

```
lymphnodesremoved:
  Min. 1st Qu.
                 Median
                           Mean 3rd Qu.
                                            Max.
  11.00
        13.25
                  16.50
                          19.00
                                  22.75
                                           42.00
lymphnodesinvaded:
  Min. 1st Qu.
                Median
                           Mean 3rd Qu.
                                            Max.
  0.000
         0.000
                  0.000
                          1.903
                                 1.750
                                          20.000
stageall:
1 2 3
         4
13 23 21 5
dfs_status:
deceased_or_recurrence
                          living_norecurrence
                                                                  NA's
                                                                     1
                                            46
```

batch:

28 34

Length Class Mode 62 character character

uncurated\_author\_metadata:
Length Class Mode
62 character character

GSE13067\_eset

DNA copy-number alterations underlie gene expression differences between microsatellite stable and unstable colorectal cancers.

# **Description**

About 15% of colorectal cancers harbor microsatellite instability (MSI). MSI-associated gene expression changes have been identified in colorectal cancers, but little overlap exists between signatures hindering an assessment of overall consistency. Little is known about the causes and downstream effects of differential gene expression. DNA microarray data on 89 MSI and 140 microsatellite-stable (MSS) colorectal cancers from this study and 58 MSI and 77 MSS cases from three published reports were randomly divided into test and training sets. MSI-associated gene expression changes were assessed for cross-study consistency using training samples and validated as MSI classifier using test samples. Differences in biological pathways were identified by functional category analysis. Causation of differential gene expression was investigated by comparison to DNA copy-number

GSE13067\_eset 13

data.MSI-associated gene expression changes in colorectal cancers were found to be highly consistent across multiple studies of primary tumors and cancer cell lines from patients of different ethnicities (P < 0.001). Clustering based on consistent changes separated additional test cases by MSI status, and classification of individual samples predicted MSI status with a sensitivity of 96% and specificity of 85%. Genes associated with immune response were up-regulated in MSI cancers, whereas genes associated with cell-cell adhesion, ion binding, and regulation of metabolism were down-regulated. Differential gene expression was shown to reflect systematic differences in DNA copy-number aberrations between MSI and MSS tumors (P < 0.001). Our results show cross-study consistency of MSI-associated gene expression changes in colorectal cancers. DNA copy-number alterations partly cause the differences in gene expression between MSI and MSS cancers.

# Usage

```
data( GSE13067_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Jorissen RN, Lipton L, Gibbs P, Chapman M et al.??DNA copy-number alterations under
 Laboratory: Jorissen and Sieber 2008
 Contact information:
 Title: DNA copy-number alterations underlie gene expression differences between microsatellite stable
 URL:
 PMIDs: 19088021
  Abstract: A 251 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
```

14 *GSE13294\_eset* 

```
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
```

varLabels: probeset gene
varMetadata: labelDescription

#### **Details**

```
assayData: 19320 features, 74 samples
Platform type: hgu133plus2
Available sample meta-data:
alt_sample_name:
   Length
              Class
                         Mode
       74 character character
sample_type:
tumor
   74
msi:
MSI MSS
 11 63
batch:
   Length
              Class
                         Mode
       74 character character
uncurated_author_metadata:
   Length
              Class
       74 character character
```

GSE13294\_eset

DNA copy-number alterations underlie gene expression differences between microsatellite stable and unstable colorectal cancers.

# Description

About 15% of colorectal cancers harbor microsatellite instability (MSI). MSI-associated gene expression changes have been identified in colorectal cancers, but little overlap exists between signatures hindering an assessment of overall consistency. Little is known about the causes and downstream effects of differential gene expression.DNA microarray data on 89 MSI and 140 microsatellite-stable (MSS) colorectal cancers from this study and 58 MSI and 77 MSS cases from three published reports were randomly divided into test and training sets. MSI-associated gene expression changes

GSE13294\_eset 15

were assessed for cross-study consistency using training samples and validated as MSI classifier using test samples. Differences in biological pathways were identified by functional category analysis. Causation of differential gene expression was investigated by comparison to DNA copy-number data.MSI-associated gene expression changes in colorectal cancers were found to be highly consistent across multiple studies of primary tumors and cancer cell lines from patients of different ethnicities (P < 0.001). Clustering based on consistent changes separated additional test cases by MSI status, and classification of individual samples predicted MSI status with a sensitivity of 96% and specificity of 85%. Genes associated with immune response were up-regulated in MSI cancers, whereas genes associated with cell-cell adhesion, ion binding, and regulation of metabolism were down-regulated. Differential gene expression was shown to reflect systematic differences in DNA copy-number aberrations between MSI and MSS tumors (P < 0.001). Our results show cross-study consistency of MSI-associated gene expression changes in colorectal cancers. DNA copy-number alterations partly cause the differences in gene expression between MSI and MSS cancers.

#### Usage

```
data( GSE13294_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Jorissen RN, Lipton L, Gibbs P, Chapman M et al. DNA copy-number alterations underli
 Laboratory: Jorissen and Sieber 2008
 Contact information:
 Title: DNA copy-number alterations underlie gene expression differences between microsatellite stable
 PMIDs: 19088021
 Abstract: A 251 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
     [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
  platform_shorttitle:
      Affymetrix HG-U133Plus2
  platform_summary:
     hgu133plus2
  platform_manufacturer:
     Affymetrix
  platform_distribution:
      commercial
  platform_accession:
     GPL570
  platform_technology:
     in situ oligonucleotide
  warnings:
     No warnings yet
```

16 GSE14095\_eset

```
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
varLabels: probeset gene
varMetadata: labelDescription
```

#### **Details**

```
assayData: 19320 features, 155 samples
Platform type: hgu133plus2
Available sample meta-data:
alt_sample_name:
   Length Class
                         Mode
      155 character character
sample_type:
tumor
  155
msi:
MSI MSS
78 77
batch:
   Length
              Class
                         Mode
      155 character character
uncurated_author_metadata:
   Length
             Class
      155 character character
```

GSE14095\_eset

Gene expression signature and response to the use of leucovorin, fluorouracil and oxaliplatin in colorectal cancer patients.

#### **Description**

FOLFOX (a combination of leucovorin, fluorouracil and oxaliplatin) has achieved substantial success in the treatment of colorectal cancer (CRC) patients. However, about half of all patients show resistance to this regimen and some develop adverse symptoms such as neurotoxicity. In order to select patients who would benefit most from this therapy, we aimed to build a predictor for the

GSE14095\_eset 17

response to FOLFOX using microarray gene expression profiles of primary CRC samples. Forty patients who underwent surgery for primary lesions were examined. All patients had metastatic or recurrent CRC and received modified FOLFOX6. Responders and nonresponders were determined according to the best observed response at the end of the first-line treatment. Gene-expression profiles of primary CRC were determined using Human Genome GeneChip arrays U133. We identified discriminating genes whose expression differed significantly between responders and nonresponders and then carried out supervised class prediction using the k-nearest-neighbour method. We identified 27 probes that were differentially expressed between responders and nonresponders at significant levels. Based on the expression of these genes, we constructed a FOLFOX response predictor with an overall accuracy of 92.5%. The sensitivity, specificity, positive and negative predictive values were 78.6%, 100%, 100% and 89.7%, respectively. The present model suggests the possibility of selecting patients who would benefit from FOLFOX therapy both in the metastatic and the adjuvant setting. To our knowledge, this is the first study to establish a prediction model for the response to FOLFOX chemotherapy based on gene expression by microarray analysis.

#### Usage

```
data( GSE14095_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Watanabe T, Kobunai T, Yamamoto Y, Matsuda K et al.??Gene expression signature and r
 Laboratory: Watanabe, Hashimoto 2008
 Contact information:
 Title: Gene expression signature and response to the use of leucovorin, fluorouracil and oxaliplatin in
 URL:
 PMIDs: 21680303
 Abstract: A 241 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
```

18 *GSE14333\_eset* 

```
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
varLabels: probeset gene
varMetadata: labelDescription
```

#### **Details**

```
assayData: 19320 features, 189 samples
Platform type: hgu133plus2
_____
Available sample meta-data:
_____
alt_sample_name:
  Length
            Class
                      Mode
     189 character character
sample_type:
tumor
  189
drug_response:
       y NA's
  n
  5
       5 179
uncurated_author_metadata:
  Length
            Class
     189 character character
```

GSE14333\_eset

Metastasis-Associated Gene Expression Changes Predict Poor Outcomes in Patients with Dukes Stage B and C Colorectal Cancer.

# Description

PURPOSE: Colorectal cancer prognosis is currently predicted from pathologic staging, providing limited discrimination for Dukes stage B and C disease. Additional markers for outcome are required to help guide therapy selection for individual patients. EXPERIMENTAL DESIGN: A multisite single-platform microarray study was done on 553 colorectal cancers. Gene expression changes were identified between stage A and D tumors (three training sets) and assessed as a prognosis signature in stage B and C tumors (independent test and external validation sets). RESULTS: One hundred twenty-eight genes showed reproducible expression changes between three sets of

GSE14333\_eset 19

stage A and D cancers. Using consistent genes, stage B and C cancers clustered into two groups resembling early-stage and metastatic tumors. A Prediction Analysis of Microarray algorithm was developed to classify individual intermediate-stage cancers into stage A-like/good prognosis or stage D-like/poor prognosis types. For stage B patients, the treatment adjusted hazard ratio for 6-year recurrence in individuals with stage D-like cancers was 10.3 (95% confidence interval, 1.3-80.0; P = 0.011). For stage C patients, the adjusted hazard ratio was 2.9 (95% confidence interval, 1.1-7.6; P = 0.016). Similar results were obtained for an external set of stage B and C patients. The prognosis signature was enriched for downregulated immune response genes and upregulated cell signaling and extracellular matrix genes. Accordingly, sparse tumor infiltration with mononuclear chronic inflammatory cells was associated with poor outcome in independent patients. CONCLUSIONS: Metastasis-associated gene expression changes can be used to refine traditional outcome prediction, providing a rational approach for tailoring treatments to subsets of patients. (Clin Cancer Res 2009;15(24):7642-51).

#### Usage

```
data( GSE14333_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Jorissen RN, Gibbs P, Christie M, Prakash S et al.??Metastasis-Associated Gene Expr
 Laboratory: Jorissen and Sieber 2008
 Contact information:
 Title: Metastasis-Associated Gene Expression Changes Predict Poor Outcomes in Patients with Dukes Stag
 PMIDs: 19996206
 Abstract: A 257 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
  platform_shorttitle:
      Affymetrix HG-U133Plus2
  platform_summary:
     hgu133plus2
  platform_manufacturer:
     Affymetrix
  platform_distribution:
      commercial
  platform_accession:
      GPL570
  platform_technology:
     in situ oligonucleotide
  warnings:
     No warnings yet
```

20 GSE14333\_eset

```
Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19320 features, 290 samples
   Platform type: hgu133plus2
   Available sample meta-data:
   _____
   alt_sample_name:
      Length Class
                          Mode
         290 character character
   sample_type:
   tumor
     290
   primarysite:
```

co re NA's 39 1

summarystage: early late NA's 44 152

0 NA's 138 152

250

N:

M: 0 1 229 61

Dstage: A B C D 44 94 91 61

summarylocation:

age\_at\_initial\_pathologic\_diagnosis: Min. 1st Qu. Median Mean 3rd Qu.

26.00 58.00 67.00 65.96 75.00 92.00

Max.

```
r NA's
  1
161 125
gender:
 f m
126 164
stageall:
  1
       3
            4 NA's
 44
      91
           61
               94
dfs_status:
deceased_or_recurrence
                         living_norecurrence
                                                               NA's
                   50
                                         176
                                                                 64
days_to_recurrence_or_death:
  Min. 1st Qu. Median
                          Mean 3rd Qu.
                                          Max.
                                                  NA's
  27.6 668.5 1154.0 1306.0 1785.0 4276.0
                                                    64
batch:
  Length
             Class
                        Mode
     290 character character
drug_treatment:
  n y NA's
172 117
preop_drug_treatment:
290
chemotherapy:
       y NA's
172 117 1
uncurated_author_metadata:
            Class
     290 character character
```

GSE16125.GPL5175\_eset Integrative approach for prioritizing cancer genes in sporadic colon cancer.

#### **Description**

The current multistep carcinogenesis models of colon cancer do not fully capture the genetic heterogeneity of the disease, which is additionally complicated by the presence of passenger and driver

genetic alterations. The aim of this study was to select in the context of this significant heterogeneity additional genes functionally related to colon cancer development. High-throughput copy number and gene expression data of 36 microsatellite stable sporadic colon cancers resected from patients of a single institution characterized for mutations in APC, KRAS, TP53 and loss of 18q were analyzed. Genes whose expression correlated with the underlying copy number pattern were selected, and their association with the above listed mutations and overall survival was evaluated. Gain of 20q was strongly associated with TP53 mutation, and overall survival with alterations on 7p, 8p, 13q, 18q, and 20q. An association with 18q loss and gain of 8q24 was also observed. New candidate genes with a potential role in colon cancer are PLCG1 on 20q, DBC1 on 8q21, and NDGR1 on 8p24. In addition, an unexpected pattern of loss and mutability was found in the region upstream of the KRAS gene. By integrating copy number alterations with gene expression and mutations in colon cancer associated genes, we have developed a strategy that identifies previously known molecular features and additional players in the molecular landscape of colon cancer. Copyright 2009 Wiley-Liss, Inc.

#### Usage

```
data( GSE16125.GPL5175_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Reid JF, Gariboldi M, Sokolova V, Capobianco P et al.??Integrative approach for price
 Laboratory: Reid, Pierotti 2009
 Contact information:
 Title: Integrative approach for prioritizing cancer genes in sporadic colon cancer.
 URL:
 PMIDs: 19672874
 Abstract: A 225 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
     [HuEx-1_0-st] Affymetrix Human Exon 1.0 ST Array [transcript (gene) versio
n]
   platform_shorttitle:
      Affymetrix HuEx-1_0-st
   platform_summary:
      huex.1.0.st.v2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL5175
   platform_technology:
      in situ oligonucleotide
   warnings:
```

```
No warnings yet

Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
featureNames: A1CF A2ML1 ... ZYX (10015 total)
varLabels: probeset gene
varMetadata: labelDescription

Details
assayData: 10015 features, 36 samples
Platform type: huex.1.0.st.v2
```

```
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)
  4 observations deleted due to missingness
records n.max n.start events median 0.95LCL 0.95UCL
  32.00 32.00 32.00 10.00
                              6.16 5.26
Available sample meta-data:
-----
alt_sample_name:
  Length
         Class
      36 character character
sample_type:
tumor
  36
summarystage:
early late NA's
   1
       27
M:
     1 NA's
  0
 14 19
Dstage:
  B C
         D NA's
         19 4
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median Mean 3rd Qu.
                                      Max.
  19.00 61.75 67.50 65.58 71.25
                                     90.00
```

```
days_to_death:
  Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
  30.0 202.5 390.0 926.2 1658.0 2970.0 4
vital_status:
deceased living NA's
  10 22
family_history:
n
36
msi:
MSS
36
gender:
f m
20 16
kras:
mutant
   20
      16
tumor_size:
  Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
 2.500 4.000 4.000 4.906 5.000 14.000
mutation_apc:
n y
8 28
stageall:
 1 2 3 4 NA's
  1 5 8 19 3
batch:
  Length
          Class Mode
     36 character character
preop_drug_treatment:
n
36
uncurated_author_metadata:
  Length Class Mode
     36 character character
```

GSE17536\_eset 25

GSE17536\_eset Experimentally derived metastasis gene expression profile predicts recurrence and death in patients with colon cancer.

#### **Description**

Staging inadequately predicts metastatic risk in patients with colon cancer. We used a gene expression profile derived from invasive, murine colon cancer cells that were highly metastatic in an immunocompetent mouse model to identify patients with colon cancer at risk of recurrence. This phase 1, exploratory biomarker study used 55 patients with colorectal cancer from Vanderbilt Medical Center (VMC) as the training dataset and 177 patients from the Moffitt Cancer Center as the independent dataset. The metastasis-associated gene expression profile developed from the mouse model was refined with comparative functional genomics in the VMC gene expression profiles to identify a 34-gene classifier associated with high risk of metastasis and death from colon cancer. A metastasis score derived from the biologically based classifier was tested in the Moffitt dataset.A high score was significantly associated with increased risk of metastasis and death from colon cancer across all pathologic stages and specifically in stage II and stage III patients. The metastasis score was shown to independently predict risk of cancer recurrence and death in univariate and multivariate models. For example, among stage III patients, a high score translated to increased relative risk of cancer recurrence (hazard ratio, 4.7; 95% confidence interval, 1.566-14.05). Furthermore, the metastasis score identified patients with stage III disease whose 5-year recurrence-free survival was >88% and for whom adjuvant chemotherapy did not increase survival time. A gene expression profile identified from an experimental model of colon cancer metastasis predicted cancer recurrence and death, independently of conventional measures, in patients with colon cancer. Copyright 2010 AGA Institute. Published by Elsevier Inc. All rights reserved.

#### Usage

```
data(GSE17536_eset)
```

```
experimentData(eset):
Experiment data
    Experimenter name: Smith JJ, Deane NG, Wu F, Merchant NB et al.??Experimentally derived metastasis gene
    Laboratory: Smith JJ,??Beauchamp RD 2009
    Contact information:
    Title: Experimentally derived metastasis gene expression profile predicts recurrence and death in pati
    URL:
    PMIDs: 19914252

Abstract: A 260 word abstract is available. Use 'abstract' method.
    Information is available on: preprocessing
    notes:
        platform_title:
            [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
        platform_shorttitle:
```

26 GSE17536\_eset

```
Affymetrix HG-U133Plus2
      platform_summary:
         hgu133plus2
      platform_manufacturer:
         Affymetrix
      platform_distribution:
         commercial
      platform_accession:
         GPL570
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19320 features, 177 samples
   Platform type: hgu133plus2
   Overall survival time-to-event summary (in years):
   Call: survfit(formula = Surv(time, cens) ~ -1)
   records
           n.max n.start events median 0.95LCL 0.95UCL
    177.00 177.00 177.00 73.00 11.08 4.86
   _____
   Available sample meta-data:
   -----
   alt_sample_name:
      Length
             Class
                           Mode
         177 character character
   sample_type:
   tumor
     177
   primarysite:
    СО
   177
   summarygrade:
```

GSE17536\_eset 27

```
high low
 27 150
G:
 1 2 3
16 134 27
summarystage:
early late NA's
  24 96 57
N:
 0 NA's
 81 96
M:
 0 1
138 39
Dstage:
 B C D NA's
 57 57 39 24
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median Mean 3rd Qu.
                                     Max.
 26.00 57.00 66.00 65.48 75.00 92.00
days_to_tumor_recurrence:
  Min. 1st Qu. Median
                       Mean 3rd Qu.
   0.0 298.8 858.9 1126.0 1734.0 4276.0
recurrence_status:
                             NA's
norecurrence recurrence
       109
               36
                              32
days_to_death:
  Min. 1st Qu. Median Mean 3rd Qu.
  27.6 683.4 1268.0 1444.0 2035.0 4276.0
vital_status:
deceased living
     73
          104
gender:
f m
81 96
```

stageall:

28 GSE17537\_eset

```
1 2 3 4
24 57 57 39
ethnicity:
    black caucasian hispanic
                                   other
        9
                151
                                      16
                            1
dfs_status:
deceased_or_recurrence
                          living_norecurrence
                                                                 NA's
                    36
                                            91
                                                                    50
days_to_recurrence_or_death:
  Min. 1st Qu. Median
                           Mean 3rd Qu.
                                            Max.
                                                    NA's
                                                      50
   27.6 751.3 1380.0
                        1519.0 2122.0
                                         4276.0
disease_specific_mortality:
122 55
batch:
  Length
              Class
                         Mode
      177 character character
preop_drug_treatment:
 n
177
uncurated_author_metadata:
   Length
              Class
                         Mode
      177 character character
```

GSE17537\_eset

Experimentally derived metastasis gene expression profile predicts recurrence and death in patients with colon cancer.

# **Description**

Staging inadequately predicts metastatic risk in patients with colon cancer. We used a gene expression profile derived from invasive, murine colon cancer cells that were highly metastatic in an immunocompetent mouse model to identify patients with colon cancer at risk of recurrence. This phase 1, exploratory biomarker study used 55 patients with colorectal cancer from Vanderbilt Medical Center (VMC) as the training dataset and 177 patients from the Moffitt Cancer Center as the independent dataset. The metastasis-associated gene expression profile developed from the mouse model was refined with comparative functional genomics in the VMC gene expression profiles to identify a 34-gene classifier associated with high risk of metastasis and death from colon cancer. A metastasis score derived from the biologically based classifier was tested in the Moffitt dataset. A

GSE17537\_eset 29

high score was significantly associated with increased risk of metastasis and death from colon cancer across all pathologic stages and specifically in stage II and stage III patients. The metastasis score was shown to independently predict risk of cancer recurrence and death in univariate and multivariate models. For example, among stage III patients, a high score translated to increased relative risk of cancer recurrence (hazard ratio, 4.7; 95% confidence interval, 1.566-14.05). Furthermore, the metastasis score identified patients with stage III disease whose 5-year recurrence-free survival was >88% and for whom adjuvant chemotherapy did not increase survival time. A gene expression profile identified from an experimental model of colon cancer metastasis predicted cancer recurrence and death, independently of conventional measures, in patients with colon cancer. Copyright 2010 AGA Institute. Published by Elsevier Inc. All rights reserved.

# Usage

```
data( GSE17537_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Smith JJ, Deane NG, Wu F, Merchant NB et al.??Experimentally derived metastasis gene
 Laboratory: Smith JJ,??Beauchamp RD 2009
 Contact information:
 Title: Experimentally derived metastasis gene expression profile predicts recurrence and death in pati
 URL:
 PMIDs: 19914252
  Abstract: A 260 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hg-u133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
```

30 GSE17537\_eset

```
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total) varLabels: probeset gene varMetadata: labelDescription
```

# **Details**

```
assayData: 19320 features, 55 samples
Platform type: hg-u133plus2
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)
        n.max n.start events median 0.95LCL 0.95UCL
records
    55
           55
                55
                       20
                               NA
                                       NA
-----
Available sample meta-data:
-----
alt_sample_name:
  Length
         Class
                     Mode
      55 character character
sample_type:
tumor
  55
primarysite:
СО
55
summarygrade:
high low NA's
  3 33 19
G:
     2
           3 NA's
  1
  1 32
           3 19
summarystage:
early late NA's
   4 36 15
N:
  0 NA's
 19 36
M:
0 1
```

```
GSE17537_eset
                                                                                     31
```

```
38 17
Dstage:
  B C D NA's
 15 19 17 4
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median
                       Mean 3rd Qu.
                                       Max.
 23.00 54.00 62.00
                        62.31 72.00
                                       94.00
days_to_tumor_recurrence:
  Min. 1st Qu. Median
                        Mean 3rd Qu.
  0.00 13.81 1243.00 988.10 1694.00 2303.00
recurrence_status:
norecurrence recurrence
         36
days_to_death:
  Min. 1st Qu. Median
                       Mean 3rd Qu.
 12.82 950.30 1506.00 1357.00 1801.00 3345.00
vital_status:
deceased living
     20
gender:
f m
29 26
stageall:
1 2 3 4
4 15 19 17
ethnicity:
   black caucasian hispanic
       4 50
dfs_status:
deceased_or_recurrence
                        living_norecurrence
                                                           NA's
days_to_recurrence_or_death:
  Min. 1st Qu. Median Mean 3rd Qu.
                                       Max.
                                               NA's
```

29.1 1505.0 1196.0 1782.0 3345.0

disease\_specific\_mortality:

n y

5

```
batch:
   Length Class Mode
     55 character character
uncurated_author_metadata:
   Length Class Mode
     55 character character
```

GSE17538.GPL570\_eset

Experimentally derived metastasis gene expression profile predicts recurrence and death in patients with colon cancer.

#### Description

Staging inadequately predicts metastatic risk in patients with colon cancer. We used a gene expression profile derived from invasive, murine colon cancer cells that were highly metastatic in an immunocompetent mouse model to identify patients with colon cancer at risk of recurrence. This phase 1, exploratory biomarker study used 55 patients with colorectal cancer from Vanderbilt Medical Center (VMC) as the training dataset and 177 patients from the Moffitt Cancer Center as the independent dataset. The metastasis-associated gene expression profile developed from the mouse model was refined with comparative functional genomics in the VMC gene expression profiles to identify a 34-gene classifier associated with high risk of metastasis and death from colon cancer. A metastasis score derived from the biologically based classifier was tested in the Moffitt dataset.A high score was significantly associated with increased risk of metastasis and death from colon cancer across all pathologic stages and specifically in stage II and stage III patients. The metastasis score was shown to independently predict risk of cancer recurrence and death in univariate and multivariate models. For example, among stage III patients, a high score translated to increased relative risk of cancer recurrence (hazard ratio, 4.7; 95% confidence interval, 1.566-14.05). Furthermore, the metastasis score identified patients with stage III disease whose 5-year recurrence-free survival was >88% and for whom adjuvant chemotherapy did not increase survival time. A gene expression profile identified from an experimental model of colon cancer metastasis predicted cancer recurrence and death, independently of conventional measures, in patients with colon cancer. Copyright 2010 AGA Institute. Published by Elsevier Inc. All rights reserved.

# Usage

```
data( GSE17538.GPL570_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
Experimenter name: Smith JJ, Deane NG, Wu F, Merchant NB et al.??Experimentally derived metastasis generated and the second s
```

Laboratory: Smith JJ,??Beauchamp RD 2009

```
Contact information:
     Title: Experimentally derived metastasis gene expression profile predicts recurrence and death in pati
     PMIDs: 19914252
     Abstract: A 260 word abstract is available. Use 'abstract' method.
     Information is available on: preprocessing
     notes:
      platform_title:
         [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
      platform_shorttitle:
         Affymetrix HG-U133Plus2
      platform_summary:
         hgu133plus2
      platform_manufacturer:
         Affymetrix
      platform_distribution:
         commercial
      platform_accession:
         GPL570
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19320 features, 232 samples
   Platform type: hgu133plus2
   Overall survival time-to-event summary (in years):
   Call: survfit(formula = Surv(time, cens) ~ -1)
   records
            n.max n.start events median 0.95LCL 0.95UCL
    232.00 232.00 232.00 93.00 11.08
                                              5.57
   Available sample meta-data:
   alt_sample_name:
      Length Class
                            Mode
```

```
232 character character
```

```
sample_type:
tumor
   232

summarygrade:
high low NA's
   30 183 19

G:
```

1 2 3 NA's 17 166 30 19

summarystage:
early late NA's
 28 132 72

N: 0 NA's 100 132

M: 0 1 176 56

Dstage:

B C D NA's 72 76 56 28

age\_at\_initial\_pathologic\_diagnosis:

Min. 1st Qu. Median Mean 3rd Qu. Max.
23.00 56.00 65.50 64.73 74.00 94.00

days\_to\_tumor\_recurrence:

Min. 1st Qu. Median Mean 3rd Qu. Max. 0.0 216.2 906.8 1093.0 1713.0 4276.0

recurrence\_status:

norecurrence recurrence NA's 145 55 32

days\_to\_death:

Min. 1st Qu. Median Mean 3rd Qu. Max. 12.82 699.70 1402.00 1423.00 1919.00 4276.00

vital\_status:
deceased living

GSE18105\_eset 35

```
93
              139
gender:
 f
110 122
stageall:
1 2 3 4
28 72 76 56
ethnicity:
   black caucasian hispanic
                                  other
       13
                201
                                     16
                            2
dfs_status:
deceased_or_recurrence
                          living_norecurrence
                                                                 NA's
                                                                   55
                                          122
days_to_recurrence_or_death:
  Min. 1st Qu. Median
                           Mean 3rd Qu.
                                           Max.
                                                   NA's
   0.0 738.6 1473.0 1428.0 1917.0
                                        4276.0
                                                     55
disease_specific_mortality:
  n
162 70
batch:
  Length
              Class
                         Mode
     232 character character
uncurated_author_metadata:
  Length
              Class
     232 character character
```

GSE18105\_eset

MUC12 mRNA expression is an independent marker of prognosis in stage II and stage III colorectal cancer.

# Description

Distant metastasis is the major cause of death in colorectal cancer (CRC) patients. To identify genes influencing the prognosis of patients with CRC, we compared gene expression in primary tumors with and without distant metastasis using an oligonucleotide microarray. We also examined the expression of the candidate gene in 100 CRC patients by quantitative real-time reverse transcription PCR and studied the relationship between its expression and the prognosis of patients with CRC. As a result, we identified MUC12 as a candidate gene involved in metastasis processes by microarray

36 GSE18105\_eset

analysis. Quantitative real-time reverse transcription PCR showed that MUC12 expression was significantly lower in cancer tissues than in adjacent normal tissues (p < 0.001). In Stages II and III CRC, patients with low expression showed worse disease-free survival (p = 0.020). Multivariate analysis disclosed that MUC12 expression status was an independent prognostic factor in Stages II and III CRC (relative risk, 8.236; 95% confidence interval, 1.702-39.849 p = 0.009). Our study revealed the prognostic value of MUC12 expression in CRC patients. Moreover, our result suggests MUC12 expression is a possible candidate gene for assessing postoperative adjuvant therapy for CRC patients.

#### Usage

```
data( GSE18105_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Matsuyama T, Ishikawa T, Mogushi K, Yoshida T et al.??MUC12 mRNA expression is an in
 Laboratory: Matsuyama, Sugihara 2009
 Contact information:
 Title: MUC12 mRNA expression is an independent marker of prognosis in stage II and stage III colorectal
 URL:
 PMIDs: 20162577
  Abstract: A 188 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
  varLabels: probeset gene
```

GSE18105\_eset 37

varMetadata: labelDescription

```
assayData: 19320 features, 111 samples
Platform type: hgu133plus2
_____
Available sample meta-data:
-----
alt_sample_name:
  Length
         Class Mode
     111 character character
sample_type:
adjacentnormal
                   tumor
          17
                       94
summarystage:
late NA's
 44 67
M:
0 1
85 26
recurrence_status:
norecurrence recurrence
        93
             18
stageall:
  4 NA's
 26 85
ethnicity:
other
 111
batch:
  Length
           Class
                     Mode
     111 character character
preop_drug_treatment:
 n
111
uncurated_author_metadata:
  Length Class Mode
```

38 GSE2109\_eset

111 character character

GSE2109\_eset

Expression Project for Oncology (expO)

# **Description**

EXpression Project for Oncology, International Genomics Consortium, www.intgen.org

## Usage

```
data( GSE2109_eset )
```

experimentData(eset):

```
Experiment data
 Experimenter name: EXpression Project for Oncology, International Genomics Consortium, www.intgen.org
 Laboratory: exp0, IGC 2005
 Contact information:
 Title: Expression Project for Oncology (exp0)
 URL:
 PMIDs: PMID unknown
 Abstract: A 8 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
```

GSE2109\_eset 39

```
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19320 features, 427 samples
   Platform type: hgu133plus2
   Available sample meta-data:
   alt_sample_name:
      Length
             Class
                          Mode
         427 character character
   sample_type:
   tumor
     427
   primarysite:
    co re
   343 84
   summarygrade:
   high low NA's
     75 270 82
   G:
          2
             3
                  4 NA's
      1
     10 260
             71
                 4 82
   summarystage:
   early late NA's
     166 177
                 84
   T:
      Length
                Class
                          Mode
         427 character character
   N:
             2
                   X NA's
         1
    187 101 59 3 77
   M:
               X NA's
      0
          1
    274 64
             9 80
```

An object of class 'AnnotatedDataFrame'

40 GSE21510\_eset

```
Dstage:
        В
             C
                  D NA's
   Α
  51 113 105
                  5 153
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median
                            Mean 3rd Qu.
                                             Max.
  25.00
          55.00
                  65.00
                           65.68
                                   75.00
                                            95.00
gender:
   f
        m NA's
207 219
             1
stageall:
  Min. 1st Qu.
                 Median
                            Mean 3rd Qu.
                                             Max.
                                                     NA's
  0.000
          2.000
                  2.000
                           2.516
                                   3.000
                                            4.000
                                                       90
ethnicity:
    black caucasian hispanic
                                   other
       23
                391
                                      12
batch:
              Class
   Length
                         Mode
      427 character character
uncurated author metadata:
   Length
              Class
      427 character character
```

GSE21510\_eset

Clinical significance of osteoprotegerin expression in human colorectal cancer.

# Description

This study aimed to identify a novel biomarker or a target of treatment for colorectal cancer (CRC). The expression profiles of cancer cells in 104 patients with CRC were examined using laser microdissection and oligonucleotide microarray analysis. Overexpression in CRC cells, especially in patients with distant metastases, was a prerequisite to select candidate genes. The mRNA expression of candidate genes was investigated by quantitative reverse transcriptase PCR (RT-PCR) in 77 patients as a validation study. We analyzed the protein expression and localization of the candidate gene by immunohistochemical study and investigated the relationship between protein expression and clinicopathologic features in 274 CRC patients. Using microarray analysis, we identified 6 candidate genes related to distant metastases in CRC patients. Among these genes, osteoprotegerin (OPG) is known to be associated with aggressiveness in several cancers through inhibition of apoptosis

GSE21510\_eset 41

via neutralization of the function of TNF-related apoptosis-inducing ligand. The mRNA expression of OPG in cancer tissues was significantly higher in patients with distant metastases than those without metastases. Overexpression of OPG protein was associated with significantly worse overall survival and relapse-free survival. Moreover, overexpression of the OPG protein was an independent risk factor for CRC recurrence. Overexpression of OPG may be a predictive biomarker of CRC recurrence and a target for treatment of this disease.??2011 AACR.

## Usage

```
data( GSE21510_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Tsukamoto S, Ishikawa T, Iida S, Ishiguro M et al.??Clinical significance of osteopo
 Laboratory: Tsukamoto, Sugihara 2010
 Contact information:
 Title: Clinical significance of osteoprotegerin expression in human colorectal cancer.
 URL:
 PMIDs: 21270110
 Abstract: A 211 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

42 GSE21815\_eset

```
assayData: 19320 features, 148 samples
Platform type: hgu133plus2
Available sample meta-data:
alt_sample_name:
  Length Class
     148 character character
sample_type:
adjacentnormal
                      tumor
           25
                       123
summarystage:
early late NA's
  19
        74
            55
M:
 0
    1
121 27
Dstage:
  В С
         D NA's
  54 47 27 20
stageall:
 0 1 2 3 4
 1 19 54 47 27
ethnicity:
other
  148
batch:
             Class
                       Mode
   Length
     148 character character
uncurated_author_metadata:
  Length
            Class
                       Mode
     148 character character
```

GSE21815\_eset 43

GSE21815\_eset

Long noncoding RNA HOTAIR regulates polycomb-dependent chromatin modification and is associated with poor prognosis in colorectal cancers.

## Description

The functional impact of recently discovered long noncoding RNAs (ncRNAs) in human cancer remains to be clarified. One long ncRNA which has attracted attention is the Hox transcript antisense intergenic RNA termed HOTAIR, a long ncRNA expressed from the developmental HOXC locus located on chromosome 12q13.13. In cooperation with Polycomb complex PRC2, the HOTAIR long ncRNA is reported to reprogram chromatin organization and promote breast cancer metastasis. In this study, we examined the status and function of HOTAIR in patients with stage IV colorectal cancer (CRC) who have liver metastases and a poor prognosis. HOTAIR expression levels were higher in cancerous tissues than in corresponding noncancerous tissues and high HOTAIR expression correlated tightly with the presence of liver metastasis. Moreover, patients with high HOTAIR expression had a relatively poorer prognosis. In a subset of 32 CRC specimens, gene set enrichment analysis using cDNA array data revealed a close correlation between expression of HOTAIR and members of the PRC2 complex (SUZ12, EZH2, and H3K27me3). Our findings suggest that HOTAIR expression is associated with a genome-wide reprogramming of PRC2 function not only in breast cancer but also in CRC, where upregulation of this long ncRNA may be a critical element in metastatic progression.

## Usage

```
data( GSE21815_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Kogo R, Shimamura T, Mimori K, Kawahara K et al.??Long noncoding RNA HOTAIR regulate
 Laboratory: Mori M,??Mimori K,??Yokobori T 2010
 Contact information:
 Title: Long noncoding RNA HOTAIR regulates polycomb-dependent chromatin modification and is associated
 URL:
 PMIDs: 21862635
 Abstract: A 201 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
     Agilent-014850 Whole Human Genome Microarray 4x44K G4112F (Probe Name vers
  platform_shorttitle:
     Agilent G4112F
  platform_summary:
     hgug4112a
  platform_manufacturer:
```

44 GSE21815\_eset

```
Agilent
      platform_distribution:
         commercial
      platform_accession:
         GPL6480
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: default
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1BG A1BG-AS1 ... ZZZ3 (19686 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19686 features, 141 samples
   Platform type: hgug4112a
   _____
   Available sample meta-data:
   _____
   alt_sample_name:
      Length
                          Mode
                Class
         141 character character
   sample_type:
   adjacentnormal
                        tumor
                           132
   summarygrade:
   high low NA's
      1
         65
             75
   G:
      1
          2
               3 NA's
     32
         33
             1 75
   summarystage:
   early late NA's
      53
         78 10
   T:
      Length
                Class
                          Mode
         141 character character
```

```
N:
   0
        1
              2
                   X NA's
  70
       34
             27
                   1
                        9
M:
   0
        1 NA's
115
       17
              9
Dstage:
   Α
        В
             С
                   D NA's
   5
       59
             49
                  17
                       11
age_at_initial_pathologic_diagnosis:
                                                       NA's
   Min. 1st Qu.
                  Median
                            Mean 3rd Qu.
                                              Max.
  32.00
          57.00
                   65.00
                            64.73
                                   74.00
                                             85.00
                                                          4
gender:
        m NA's
   f
       78
  59
             4
stageall:
                                                      NA's
   Min. 1st Qu.
                  Median
                            Mean 3rd Qu.
                                              Max.
  0.000
          2.000
                   3.000
                            2.466
                                    3.000
                                             4.000
                                                         10
ethnicity:
other
  141
preop_drug_treatment:
141
uncurated_author_metadata:
               Class
   Length
                          Mode
      141 character character
```

GSE24549.GPL5175\_eset Transcriptome instability in colorectal cancer identified by exon microarray analyses: Associations with splicing factor expression levels and patient survival.

# **Description**

Colorectal cancer (CRC) is a heterogeneous disease that, on the molecular level, can be characterized by inherent genomic instabilities; chromosome instability and microsatellite instability. In the

present study we analyze genome-wide disruption of pre-mRNA splicing, and propose transcriptome instability as a characteristic that is analogous to genomic instability on the transcriptome level. Exon microarray profiles from two independent series including a total of 160 CRCs were investigated for their relative amounts of exon usage differences. Each exon in each sample was assigned an alternative splicing score calculated by the FIRMA algorithm. Amounts of deviating exon usage per sample were derived from exons with extreme splicing scores. There was great heterogeneity within both series in terms of sample-wise amounts of deviating exon usage. This was strongly associated with the expression levels of approximately half of 280 splicing factors (54% and 48% of splicing factors were significantly correlated to deviating exon usage amounts in the two series). Samples with high or low amounts of deviating exon usage, associated with overall transcriptome instability, were almost completely separated into their respective groups by hierarchical clustering analysis of splicing factor expression levels in both sample series. Samples showing a preferential tendency towards deviating exon skipping or inclusion were associated with skewed transcriptome instability. There were significant associations between transcriptome instability and reduced patient survival in both sample series. In the test series, patients with skewed transcriptome instability showed the strongest prognostic association (P = 0.001), while a combination of the two characteristics showed the strongest association with poor survival in the validation series (P = 0.03). We have described transcriptome instability as a characteristic of CRC. This transcriptome instability has associations with splicing factor expression levels and poor patient survival.

# Usage

```
data( GSE24549.GPL5175_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Sveen A,????esen TH,??Rognum TO,??Lothe RA,??Skotheim RI. Transcriptome instabilit
 Laboratory: Sveen A,????esen TH,??Rognum TO,??Lothe RA,??Skotheim RI 2011
 Contact information:
 Title: Transcriptome instability in colorectal cancer identified by exon microarray analyses: Associa-
  URL:
 PMIDs: 21619627
  Abstract: A 282 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HuEx-1_0-st] Affymetrix Human Exon 1.0 ST Array [transcript (gene) versio
n
   platform_shorttitle:
      Affymetrix HuEx-1_0-st
   platform_summary:
      huex.1.0.st.v2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
```

```
platform_accession:
    GPL5175
platform_technology:
    in situ oligonucleotide
warnings:
    No warnings yet

Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
    featureNames: A1CF A2ML1 ... ZYX (10015 total)
    varLabels: probeset gene
    varMetadata: labelDescription
```

```
assayData: 10015 features, 83 samples
Platform type: huex.1.0.st.v2
Available sample meta-data:
-----
alt_sample_name:
  Length Class
                      Mode
      83 character character
sample_type:
tumor
  83
summarystage:
late NA's
 37 46
M:
0
83
Dstage:
В С
46 37
msi:
MSI MSS NA's
  7 75
stageall:
2 3
```

```
46 37
dfs_status:
deceased_or_recurrence
                          living_norecurrence
days_to_recurrence_or_death:
  Min. 1st Qu. Median
                           Mean 3rd Qu.
                                            Max.
          441.6 2394.0 2113.0 3650.0
                                         3650.0
batch:
              Class
                         Mode
   Length
       83 character character
drug_treatment:
83
preop_drug_treatment:
n
83
chemotherapy:
n y
46 37
uncurated_author_metadata:
  Length
              Class
                         Mode
       83 character character
```

GSE24550.GPL5175\_eset Transcriptome instability in colorectal cancer identified by exon microarray analyses: Associations with splicing factor expression levels and patient survival.

## **Description**

Colorectal cancer (CRC) is a heterogeneous disease that, on the molecular level, can be characterized by inherent genomic instabilities; chromosome instability and microsatellite instability. In the present study we analyze genome-wide disruption of pre-mRNA splicing, and propose transcriptome instability as a characteristic that is analogous to genomic instability on the transcriptome level. Exon microarray profiles from two independent series including a total of 160 CRCs were investigated for their relative amounts of exon usage differences. Each exon in each sample was assigned an alternative splicing score calculated by the FIRMA algorithm. Amounts of deviating exon usage per sample were derived from exons with extreme splicing scores. There was great heterogeneity within both series in terms of sample-wise amounts of deviating exon usage. This was

strongly associated with the expression levels of approximately half of 280 splicing factors (54% and 48% of splicing factors were significantly correlated to deviating exon usage amounts in the two series). Samples with high or low amounts of deviating exon usage, associated with overall transcriptome instability, were almost completely separated into their respective groups by hierarchical clustering analysis of splicing factor expression levels in both sample series. Samples showing a preferential tendency towards deviating exon skipping or inclusion were associated with skewed transcriptome instability. There were significant associations between transcriptome instability and reduced patient survival in both sample series. In the test series, patients with skewed transcriptome instability showed the strongest prognostic association (P = 0.001), while a combination of the two characteristics showed the strongest association with poor survival in the validation series (P = 0.03). We have described transcriptome instability as a characteristic of CRC. This transcriptome instability has associations with splicing factor expression levels and poor patient survival.

## Usage

```
data( GSE24550.GPL5175_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Sveen A, Agesen TH, Nesbakken A, Rognum TO et al.??Transcriptome instability in colo
 Laboratory: Sveen A,????esen TH,??Rognum TO,??Lothe RA,??Skotheim RI 2011
 Contact information:
 Title: Transcriptome instability in colorectal cancer identified by exon microarray analyses: Associa
 URL:
 PMIDs: 21619627
  Abstract: A 282 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
     [HuEx-1_0-st] Affymetrix Human Exon 1.0 ST Array [transcript (gene) versio
nΤ
   platform_shorttitle:
      Affymetrix HuEx-1_0-st
   platform_summary:
      huex.1.0.st.v2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL5175
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
```

Preprocessing: default featureData(eset): An object of class 'AnnotatedDataFrame' featureNames: A1CF A2ML1 ... ZYX (10015 total) varLabels: probeset gene

varMetadata: labelDescription

# **Details**

assayData: 10015 features, 90 samples Platform type: huex.1.0.st.v2 Available sample meta-data: \_\_\_\_\_ alt\_sample\_name: Length Class Mode 90 character character sample\_type: adjacentnormal tumor 13 77 summarystage: late NA's 33 57 M: 0 NA's 77 13 Dstage: B C NA's 44 33 13 msi: MSI MSS NA's 14 51 25 stageall: 3 NA's 2 44 33 13 dfs\_status: deceased\_or\_recurrence living\_norecurrence NA's 19 58 13 days\_to\_recurrence\_or\_death:

GSE2630\_eset 51

```
Min. 1st Qu. Median
                            Mean 3rd Qu.
                                                     NA's
  62.05 1007.00 1219.00 1193.00 1449.00 1814.00
                                                       13
batch:
   Length
              Class
                          Mode
       90 character character
drug_treatment:
   n
        y NA's
  44
       33
           13
preop_drug_treatment:
n
90
chemotherapy:
        y NA's
   n
  44
       33
           13
uncurated_author_metadata:
   Length
              Class
                          Mode
       90 character character
```

GSE2630\_eset

A gene signature of 8 genes could identify the risk of recurrence and progression in Dukes' B colon cancer patients.

## Description

The benefit of postoperative adjuvant chemotherapy in patients with Dukes' B colorectal cancer is still uncertain and its routine use is not recommended. The five-year relapse rate is approximately 25-40% and the identification of patients at high risk of recurrence would represent an important strategy for the use of adjuvant chemotherapy. We retrospectively analyzed gene expression profiles in frozen tumor specimens from patients with Dukes' B colorectal cancer by using high density oligonucleotide microarrays. Our results show a subset of 48 genes differentially expressed with an associated probability <0.001 in the t-test. Another statistical procedure based on the Fisher criterion resulted in 11 genes able to separate both groups. We selected the 8 genes present in both subsets. The differential expression of five genes (CHD2, RPS5, ZNF148, BRI3 and MGC23401) in colon cancer progression was confirmed by real-time PCR in an independent set of patients of Dukes' B and C stages.

## Usage

```
data( GSE2630_eset )
```

52 GSE2630\_eset

## **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Bandr??E,??Malumbres R,??Cubedo E,??Sola J,??Garc??Foncillas J,??Labarga A. A gene
 Laboratory: Bandr??E,??Malumbres R,??Cubedo E,??Sola J,??Garc??Foncillas J,??Labarga A 2005
 Contact information:
 Title: A gene signature of 8 genes could identify the risk of recurrence and progression in Dukes' B col
 URL:
 PMIDs: 17390049
  Abstract: A 151 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      Human 19K oligo array
   platform_shorttitle:
      NA
   platform_summary:
      human 19k oligo array
   platform_manufacturer:
   platform_distribution:
      non-commercial
   platform_accession:
      GPL2006
   platform_technology:
      spotted oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG-AS1 A1CF ... ZZZ3 (12982 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

```
assayData: 12982 features, 16 samples
Platform type: human 19k oligo array
------
Available sample meta-data:
-----
alt_sample_name:
Length Class Mode
```

GSE2630\_eset 53

```
sample_type:
tumor
  16
primarysite:
СО
16
summarystage:
early
  16
T:
 2 3
 6 10
N:
 0
16
M:
 0
16
Dstage:
В
16
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median Mean 3rd Qu.
                                        Max.
  42.00 59.50 67.00 64.38 72.00 81.00
recurrence_status:
norecurrence recurrence
summarylocation:
1 r
12 4
gender:
f m
 5 11
stageall:
```

1 2

16 character character

```
6 10
drug_treatment:
n
16
preop_drug_treatment:
16
ancillary:
n
16
chemotherapy:
16
moltherapy:
n
16
uncurated_author_metadata:
   Length
               Class
       16 character character
```

GSE26682.GPL570\_eset

MRE11 deficiency increases sensitivity to poly(ADP-ribose) polymerase inhibition in microsatellite unstable colorectal cancers.

# Description

Microsatellite instability (MSI) is displayed by approximately 15% of colorectal cancers (CRC). Defective DNA mismatch repair generates mutations at repetitive DNA sequences such as those located in the double strand break (DSB) repair gene MRE11. We assessed the mutational status of MRE11 in a panel of 17 CRC cell lines and 46 primary tumors and found a strong correlation with MSI status in both cell lines and tumors. Therefore, we hypothesized that deficiency in MRE11 may sensitize CRC cells to poly(ADP-ribose) polymerase (PARP-1) inhibition based on the concept of synthetic lethality. We further assessed the activity of the PARP-1 inhibitor, ABT-888, in CRC cell lines and observed preferential cytotoxicity in those MSI cell lines harboring mutations in MRE11 compared with both wild-type cell lines and microsatellite stable (MSS) cell lines. A significant correlation between MRE11 expression levels and cytotoxicity to ABT-888 at 10 ??M was observed (R?? = 0.915, P < 0.001). Using two experimental approaches, including short hairpin RNA knocking down MRE11 in the wild-type and MSS cell line SW-480 and a second cell line model transfected with mutant MRE11, we experimentally tried to confirm the role of MRE11 in conferring sensitivity to PARP-1 inhibition. Both models led to changes in proliferation in response

to ABT-888 at different concentrations, and a drug-response effect was not observed, suggesting a possible contribution of additional genes. We conclude that MSI colorectal tumors deficient in DSB repair secondary to mutation in MRE11 show a higher sensitivity to PARP-1 inhibition. Further clinical investigation of PARP-1 inhibitors is warranted in MSI CRCs.

### **Usage**

```
data( GSE26682.GPL570_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Vilar E,??Gruber SB,??Rennert G,??Bartnik CM,??Stenzel SL,??Iniesta MD,??Raskin L,
  Laboratory: Vilar E,??Morgan MA 2011
  Contact information:
 Title: MRE11 deficiency increases sensitivity to poly(ADP-ribose) polymerase inhibition in microsatel
  URL:
  PMIDs: 21300766
  Abstract: A 257 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1CF A2M ... ZZZ3 (11838 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

## **Details**

```
assayData: 11838 features, 156 samples
Platform type: hgu133plus2
Available sample meta-data:
alt_sample_name:
  Length
              Class
                         Mode
      156 character character
sample_type:
tumor
  156
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median
                           Mean 3rd Qu.
                                            Max.
  45.00
          65.75
                73.00
                          71.50
                                  78.00
                                          92.00
msi:
MSI MSS NA's
  15 128
            13
gender:
f m
75 81
batch:
  Length
              Class
                         Mode
      156 character character
uncurated_author_metadata:
  Length
             Class
      156 character character
```

GSE26682.GPL96\_eset

MRE11 deficiency increases sensitivity to poly(ADP-ribose) polymerase inhibition in microsatellite unstable colorectal cancers.

# **Description**

Microsatellite instability (MSI) is displayed by approximately 15% of colorectal cancers (CRC). Defective DNA mismatch repair generates mutations at repetitive DNA sequences such as those located in the double strand break (DSB) repair gene MRE11. We assessed the mutational status of MRE11 in a panel of 17 CRC cell lines and 46 primary tumors and found a strong correlation with

MSI status in both cell lines and tumors. Therefore, we hypothesized that deficiency in MRE11 may sensitize CRC cells to poly(ADP-ribose) polymerase (PARP-1) inhibition based on the concept of synthetic lethality. We further assessed the activity of the PARP-1 inhibitor, ABT-888, in CRC cell lines and observed preferential cytotoxicity in those MSI cell lines harboring mutations in MRE11 compared with both wild-type cell lines and microsatellite stable (MSS) cell lines. A significant correlation between MRE11 expression levels and cytotoxicity to ABT-888 at 10 ??M was observed (R?? = 0.915, P < 0.001). Using two experimental approaches, including short hairpin RNA knocking down MRE11 in the wild-type and MSS cell line SW-480 and a second cell line model transfected with mutant MRE11, we experimentally tried to confirm the role of MRE11 in conferring sensitivity to PARP-1 inhibition. Both models led to changes in proliferation in response to ABT-888 at different concentrations, and a drug-response effect was not observed, suggesting a possible contribution of additional genes. We conclude that MSI colorectal tumors deficient in DSB repair secondary to mutation in MRE11 show a higher sensitivity to PARP-1 inhibition. Further clinical investigation of PARP-1 inhibitors is warranted in MSI CRCs.

# Usage

```
data( GSE26682.GPL96_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Vilar E,??Gruber SB,??Rennert G,??Bartnik CM,??Stenzel SL,??Iniesta MD,??Raskin L,
 Laboratory: Vilar E,??Morgan MA 2011
 Contact information:
 Title: MRE11 deficiency increases sensitivity to poly(ADP-ribose) polymerase inhibition in microsatel
 URL:
 PMIDs: 21300766
 Abstract: A 257 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133A] Affymetrix Human Genome U133A Array
  platform_shorttitle:
     Affymetrix HG-U133A
  platform_summary:
     hgu133a
  platform_manufacturer:
     Affymetrix
  platform_distribution:
      commercial
  platform_accession:
     GPL96
  platform_technology:
      in situ oligonucleotide
  warnings:
     No warnings yet
```

```
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
featureNames: A1CF A2M ... ZZZ3 (12986 total)
varLabels: probeset gene
varMetadata: labelDescription
```

```
assayData: 12986 features, 155 samples
Platform type: hgu133a
Available sample meta-data:
-----
alt_sample_name:
  Length
         Class
     155 character character
sample_type:
tumor
 155
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median
                       Mean 3rd Qu.
                                       Max.
 21.00 66.50 75.00 72.61 80.50
                                      94.00
msi:
MSI MSS NA's
 17 123 15
gender:
f m
69 86
batch:
  Length
            Class
                      Mode
     155 character character
uncurated_author_metadata:
  Length
          Class
                      Mode
     155 character character
```

GSE26906\_eset 59

| GSE26906_eset | Expression Profiles in Stage II Colon Cancer According to APC Gene Status. |
|---------------|--|
|               |  |

## **Description**

Colorectal cancer is one of the most common cancers in the world. Histoclinical staging is efficient, but combination with molecular markers may improve the classification of stage II cancers. Several tumor-suppressor genes have been associated with colorectal cancer, and the most frequent allelic losses have been extensively studied for their prognosis effect, but the results remain controversial. In a previous study, we found a possible influence of the chromosome 5 status in the development of liver metastases in stage II colon cancers. We have here investigated the role of the APC gene, located in chromosome arm 5q, in a series of 183 colon adenocarcinomas through a combined analysis of gene expression, mutation, allelic loss and promoter methylation, and metastasis occurrence. Point mutations were found in 73% of cases and allelic losses were found in 39%; 59% of tumors presented with a biallelic inactivation, with a very strong interdependence of the two APC hits ( $P = 2.1 \times 10(-9)$ ). No association was found between expression, number and type of APC alterations, and metastatic evolution. Our results show that the determination of APC status cannot help in the prediction of metastasis and cannot be used to subclassify stage II colon cancers.

## Usage

```
data( GSE26906_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Birnbaum DJ, Laibe S, Ferrari A, Lagarde A et al.??Expression Profiles in Stage II C
 Laboratory: Olschwang S 2011
 Contact information:
 Title: Expression Profiles in Stage II Colon Cancer According to APC Gene Status.
 URL:
 PMIDs: 22496922
  Abstract: A 199 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
```

60 GSE26906\_eset

```
commercial
platform_accession:
    GPL570
platform_technology:
    in situ oligonucleotide
warnings:
    No warnings yet

Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
    featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
    varLabels: probeset gene
    varMetadata: labelDescription
```

```
assayData: 19320 features, 90 samples
Platform type: hgu133plus2
_____
Available sample meta-data:
_____
alt_sample_name:
  Length Class
                     Mode
      90 character character
sample_type:
tumor
  90
primarysite:
СО
90
N:
0
90
M:
0
90
Dstage:
В
90
age_at_initial_pathologic_diagnosis:
```

GSE26906\_eset 61

```
Min. 1st Qu. Median
                        Mean 3rd Qu.
                                          Max.
  31.00 58.50 69.50 66.52 75.00
                                         94.00
recurrence_status:
norecurrence recurrence
         69
               21
msi:
MSS
 90
summarylocation:
1 r
65 25
gender:
 f \quad \mathsf{m}
47 43
mutation_apc:
n y
22 68
stageall:
 2
90
batch:
             Class
                        Mode
  Length
      90 character character
drug_treatment:
 n
90
preop_drug_treatment:
90
chemotherapy:
 n
90
uncurated_author_metadata:
  Length
            Class
      90 character character
```

62 GSE27544\_eset

GSE27544\_eset

Genome-wide profiling characterizes CRCs with genetic instability
and specific routes to HLA class I loss and immunoescape

## **Description**

Bernal M,??Garc??Alcalde F,??Concha ????Blanco A,??Garrido F,??Ruiz-Cabello F

# Usage

```
data( GSE27544_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Bernal M,??Garc??Alcalde F,??Concha ????Blanco A,??Garrido F,??Ruiz-Cabello F
 Laboratory: Bernal M,??Garc??Alcalde F,??Concha ????Blanco A,??Garrido F,??Ruiz-Cabello F 2011
 Contact information:
 Title: Genome-wide profiling characterizes CRCs with genetic instability and specific routes to HLA cl
 URL:
  PMIDs: PMID unknown
  Abstract: A 7 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HT_HG-U133_Plus_PM] Affymetrix HT HG-U133+ PM Array Plate
   platform_shorttitle:
      Affymetrix HT HG-U133+ PM
   platform_summary:
      hthgu133pluspm
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL13158
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1CF ... ZZZ3 (20741 total)
```

GSE28702\_eset 63

varLabels: probeset gene
varMetadata: labelDescription

### **Details**

```
assayData: 20741 features, 22 samples
Platform type: hthgu133pluspm
_____
Available sample meta-data:
alt_sample_name:
  Length
            Class
                        Mode
      22 character character
sample_type:
tumor
   22
msi:
MSI MSS
  8 14
batch:
2009-10-07
       22
uncurated_author_metadata:
  Length
             Class
                        Mode
      22 character character
```

GSE28702\_eset

Potential responders to FOLFOX therapy for colorectal cancer by Random Forests analysis.

# **Description**

Molecular characterisation using gene-expression profiling will undoubtedly improve the prediction of treatment responses, and ultimately, the clinical outcome of cancer patients. To establish the procedures to identify responders to FOLFOX therapy, 83 colorectal cancer (CRC) patients including 42 responders and 41 non-responders were divided into training (54 patients) and test (29 patients) sets. Using Random Forests (RF) algorithm in the training set, predictor genes for FOLFOX therapy were identified, which were applied to test samples and sensitivity, specificity, and out-of-bag classification accuracy were calculated. In the training set, 22 of 27 responders (81.4% sensitivity) and 23 of 27 non-responders (85.1% specificity) were correctly classified. To improve the prediction

64 GSE28702\_eset

model, we removed the outliers determined by RF, and the model could correctly classify 21 of 23 responders (91.3%) and 22 of 23 non-responders (95.6%) in the training set, and 80.0% sensitivity and 92.8% specificity, with an accuracy of 69.2% in 29 independent test samples.Random Forests on gene-expression data for CRC patients was effectively able to stratify responders to FOLFOX therapy with high accuracy, and use of pharmacogenomics in anticancer therapy is the first step in planning personalised therapy.

## Usage

```
data( GSE28702_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Tsuji S, Midorikawa Y, Takahashi T, Yagi K et al.??Potential responders to FOLFOX th
 Laboratory: Tsuji 2011
 Contact information:
 Title: Potential responders to FOLFOX therapy for colorectal cancer by Random Forests analysis.
 URL:
 PMIDs: 22095227
  Abstract: A 185 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

GSE28702\_eset 65

```
assayData: 19320 features, 83 samples
Platform type: hgu133plus2
Available sample meta-data:
alt_sample_name:
          Class
  Length
      83 character character
sample_type:
metastatic
               tumor
       27
                  56
primarysite:
 co re NA's
  33
      23 27
summarygrade:
high low
  5 78
G:
1 2 3
62 16 5
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median Mean 3rd Qu.
                                         Max.
  31.00 55.00 65.00 62.96 71.00
                                        84.00
location:
  Length
             Class
                        Mode
      83 character character
summarylocation:
      r NA's
  1
 41
     15
          27
gender:
f m
29 54
drug_name:
mfolfox6
     83
drug_treatment:
```

GSE3294\_eset

```
У
83
drug_response:
n y
41 42
preop_drug_treatment:
83
folfox:
У
83
leucovorin:
83
platin:
83
chemotherapy:
У
83
uncurated_author_metadata:
   Length
              Class
                          Mode
       83 character character
```

GSE3294\_eset

Comparative study of gene expression by cDNA microarray in human colorectal cancer tissues and normal mucosa.

## **Description**

The causative molecular pathways underlying the pathogenesis of colorectal cancer (CRC) need to be better characterized. The purpose of our study was to better understand the genetic mechanism of oncogenesis for human colorectal cancer and to identify new potential tumor markers of use in clinical practice. We used cDNA microarrays to compare gene expression profiles of colorectal biopsies from 25 CRC patients and 13 normal mucosa from adjacent non-cancerous tissues. Findings were validated by real-time PCR; in addition, western blotting and immunochemistry analysis were carried out as further confirmation of differential expression at a protein level. Comparing cancerous tissues with normal colonic mucosa we identified 584 known genes differentially expressed to a significant degree (p<0.001). Many of the transcripts that were more abundant in tumors than

GSE3294\_eset 67

in non-neoplastic tissues appear to reflect important events for colon carcinogenesis. For example, a significant number of these genes serve as apoptotic inhibitors (e.g. BFAR, BIRC1, BIRC6). Furthermore, we observed the simultaneous up-regulation of HLA-E and the down-regulation of beta2-microglobulin; these genes strongly support a potential tumor escape strategy from immune surveillance in colon cancer tissues. Our study provides new gene candidates in the pathogenesis of human CRC disease. From our results we hypothesize that CRC cells escape immune surveillance through a specific gene expression alteration; moreover, over-expression of several survival genes seems to confer a more anti-apoptotic phenotype. These genes are involved in pathways not previously implicated in CRC pathogenesis and they may provide new targets for therapy.

# Usage

```
data( GSE3294_eset )
```

```
experimentData(eset):
Experiment data
 Experimenter name: Bianchini M, Levy E, Zucchini C, Pinski V et al.??Comparative study of gene expressi
 Laboratory: Bianchini 2005
 Contact information:
 Title: Comparative study of gene expression by cDNA microarray in human colorectal cancer tissues and r
 URL:
 PMIDs: 16773188
  Abstract: A 245 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      UHN SS-Human 19Kv7
   platform_shorttitle:
      UHN SS-Human 19Kv7
   platform_summary:
      uhn ss-human 19kv7
   platform manufacturer:
   platform_distribution:
      commercial
   platform_accession:
      GPL2829
   platform_technology:
      spotted DNA/cDNA
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: AA001103 AA001104 ... Z45302 (15437 total)
```

GSE3294\_eset

varLabels: probeset gene
varMetadata: labelDescription

```
assayData: 15437 features, 24 samples
Platform type: uhn ss-human 19kv7
_____
Available sample meta-data:
alt_sample_name:
         Class Mode
  Length
      24 character character
sample_type:
tumor
  24
primarysite:
co re
17 7
summarygrade:
high low
  3 21
G:
1 2 3
10 11 3
summarystage:
early
  24
T:
1 2 3
3 1 20
N:
0 1 2
10 9 5
M:
24
Dstage:
```

GSE33113\_eset 69

```
B C
6 18
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median
                           Mean 3rd Qu.
                                            Max.
  35.00 67.25
                  76.50
                          71.46
                                  78.75
                                           89.00
summarylocation:
1 r
12 12
gender:
f m
8 16
stageall:
1 2 3
3 5 16
uncurated_author_metadata:
  Length
              Class
                         Mode
       24 character character
```

GSE33113\_eset

Mutations in the Ras-Raf Axis underlie the prognostic value of CD133 in colorectal cancer.

## **Description**

High expression of cancer stem cell (CSC) marker CD133 has been used as a predictor for prognosis in colorectal cancer (CRC), suggesting that enumeration of CSCs, using CD133, is predictive for disease progression. However, we showed recently that both CD133 mRNA and protein are not downregulated during differentiation of colon CSCs, pointing to an alternative reason for the prognostic value of CD133. We therefore set out to delineate the relation between CD133 expression and prognosis. A CRC patient series was studied for expression of CD133 and other CSC markers by microarray and quantitative PCR analysis. In addition, several common mutations were analyzed to determine the relation with CD133 expression.CD133 mRNA expression predicted relapse-free survival in our patient series, whereas several other CSC markers could not. Moreover, no correlation was found between expression of other CSC markers and CD133. Interestingly, high CD133 expression was related to mutations in K-Ras and B-Raf, and inhibition of mutant K-Ras or downstream mitogen-activated protein kinase kinase (MEK) signaling decreases CD133 expression. In addition, an activated K-Ras gene expression signature could predict CD133 expression in our patient set as well as data sets of other tumor types.CD133 expression is upregulated in CRC tumors that have a hyperactivated Ras-Raf-MEK-ERK pathway and is therefore related to mutations in K-Ras or B-Raf. As mutations in either gene have been related to poor prognosis, we conclude that CD133 expression is not indicative for CSC numbers but rather related to the mutation or activity status of the Ras-Raf pathway.

70 GSE33113\_eset

## Usage

```
data( GSE33113_eset )
```

## **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: de Sousa E Melo F, Colak S, Buikhuisen J, Koster J et al.?? Methylation of cancer-ste
 Laboratory: Medema JP,??Tanis PJ 2011
  Contact information:
 Title: Mutations in the Ras-Raf Axis underlie the prognostic value of CD133 in colorectal cancer.
 PMIDs: 22496204
  Abstract: A 247 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

## **Details**

GSE33113\_eset 71

```
alt_sample_name:
  Length Class
                     Mode
      96 character character
sample_type:
adjacentnormal
                    tumor
                      90
N:
  0 NA's
 90 6
M:
  0 NA's
 90 6
Dstage:
  B NA's
 90 6
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median Mean 3rd Qu.
                                     Max. NA's
 35.00 60.25 73.50 70.02 79.00 95.00 2
gender:
  f m NA's
 49 45 2
stageall:
  2 NA's
 90 6
days_to_recurrence_or_death:
  Min. 1st Qu. Median Mean 3rd Qu. Max.
                                             NA's
  50.0 508.8 1180.0 1234.0 1633.0 3599.0
batch:
  Length
            Class
                      Mode
      96 character character
uncurated_author_metadata:
  Length Class
      96 character character
```

\_\_\_\_\_

72 GSE39582\_eset

GSE39582\_eset Gene expression classification of colon cancer into molecular subtypes: characterization, validation, and prognostic value.

## **Description**

Colon cancer (CC) pathological staging fails to accurately predict recurrence, and to date, no gene expression signature has proven reliable for prognosis stratification in clinical practice, perhaps because CC is a heterogeneous disease. The aim of this study was to establish a comprehensive molecular classification of CC based on mRNA expression profile analyses. Fresh-frozen primary tumor samples from a large multicenter cohort of 750 patients with stage I to IV CC who underwent surgery between 1987 and 2007 in seven centers were characterized for common DNA alterations, including BRAF, KRAS, and TP53 mutations, CpG island methylator phenotype, mismatch repair status, and chromosomal instability status, and were screened with whole genome and transcriptome arrays. 566 samples fulfilled RNA quality requirements. Unsupervised consensus hierarchical clustering applied to gene expression data from a discovery subset of 443 CC samples identified six molecular subtypes. These subtypes were associated with distinct clinicopathological characteristics, molecular alterations, specific enrichments of supervised gene expression signatures (stem cell phenotype-like, normal-like, serrated CC phenotype-like), and deregulated signaling pathways. Based on their main biological characteristics, we distinguished a deficient mismatch repair subtype, a KRAS mutant subtype, a cancer stem cell subtype, and three chromosomal instability subtypes, including one associated with down-regulated immune pathways, one with up-regulation of the Wnt pathway, and one displaying a normal-like gene expression profile. The classification was validated in the remaining 123 samples plus an independent set of 1,058 CC samples, including eight public datasets. Furthermore, prognosis was analyzed in the subset of stage II-III CC samples. The subtypes C4 and C6, but not the subtypes C1, C2, C3, and C5, were independently associated with shorter relapse-free survival, even after adjusting for age, sex, stage, and the emerging prognostic classifier Oncotype DX Colon Cancer Assay recurrence score (hazard ratio 1.5, 95% CI 1.1-2.1, p???=???0.0097). However, a limitation of this study is that information on tumor grade and number of nodes examined was not available. We describe the first, to our knowledge, robust transcriptome-based classification of CC that improves the current disease stratification based on clinicopathological variables and common DNA markers. The biological relevance of these subtypes is illustrated by significant differences in prognosis. This analysis provides possibilities for improving prognostic models and therapeutic strategies. In conclusion, we report a new classification of CC into six molecular subtypes that arise through distinct biological pathways.

## Usage

```
data( GSE39582_eset )
```

```
experimentData(eset):
Experiment data
Experimenter name: Marisa L, de Reyni??A, Duval A, Selves J et al.??Gene expression classification of c
Laboratory: Marisa, Boige 2012
Contact information:
```

GSE39582\_eset 73

```
Title: Gene expression classification of colon cancer into molecular subtypes: characterization, valid
     PMIDs: 23700391
     Abstract: A 384 word abstract is available. Use 'abstract' method.
     Information is available on: preprocessing
     notes:
      platform_title:
         [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
      platform_shorttitle:
         Affymetrix HG-U133Plus2
      platform_summary:
         hgu133plus2
      platform_manufacturer:
         Affymetrix
      platform_distribution:
         commercial
      platform_accession:
         GPL570
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 19320 features, 566 samples
   Platform type: hgu133plus2
   -----
   Available sample meta-data:
   alt_sample_name:
```

Length

sample\_type:

primarysite:

tumor 566 Class

566 character character

Mode

74  $GSE39582\_eset$ 

```
СО
566
summarystage:
early late NA's
  37 265 264
N:
0 NA's
301 265
M:
 0 1
506 60
Dstage:
B C D NA's
264 205 60 37
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
  22.0 59.0 68.0 66.9 76.0
                                  97.0 1
msi:
MSI MSS NA's
75 444 47
location:
 distal proximal
  342 224
summarylocation:
l r
342 224
gender:
f m
256 310
kras:
       wt NA's
mutant
  217
        328
            21
braf:
mutant
       wt
           NA's
  51
           54
```

461

stageall:

GSE3964\_eset 75

```
2
              3
     33 264 205
                 60
dfs_status:
deceased_or_recurrence
                           living_norecurrence
                                                                   NA's
                    177
                                            380
                                                                      9
days_to_recurrence_or_death:
   Min. 1st Qu. Median
                            Mean 3rd Qu.
                                             Max.
                                                     NA's
      0
            420
                    1290
                            1461
                                    2250
                                             6030
                                                         6
batch:
   Length
              Class
                          Mode
      566 character character
drug_treatment:
        y NA's
   n
316 233
           17
preop_drug_treatment:
566
chemotherapy:
        y NA's
316
     233
            17
uncurated_author_metadata:
              Class
   Length
                          Mode
      566 character character
```

GSE3964\_eset

Deciphering cellular states of innate tumor drug responses.

#### **Description**

The molecular mechanisms underlying innate tumor drug resistance, a major obstacle to successful cancer therapy, remain poorly understood. In colorectal cancer (CRC), molecular studies have focused on drug-selected tumor cell lines or individual candidate genes using samples derived from patients already treated with drugs, so that very little data are available prior to drug treatment. Transcriptional profiles of clinical samples collected from CRC patients prior to their exposure to a combined chemotherapy of folinic acid, 5-fluorouracil and irinotecan were established using microarrays. Vigilant experimental design, power simulations and robust statistics were used to restrain the rates of false negative and false positive hybridizations, allowing successful discrimination between drug resistance and sensitivity states with restricted sampling. A list of 679 genes was established that intrinsically differentiates, for the first time prior to drug exposure, subsequently diagnosed chemo-sensitive and resistant patients. Independent biological validation performed through

76 GSE3964\_eset

quantitative PCR confirmed the expression pattern on two additional patients. Careful annotation of interconnected functional networks provided a unique representation of the cellular states underlying drug responses. Molecular interaction networks are described that provide a solid foundation on which to anchor working hypotheses about mechanisms underlying in vivo innate tumor drug responses. These broad-spectrum cellular signatures represent a starting point from which by-pass chemotherapy schemes, targeting simultaneously several of the molecular mechanisms involved, may be developed for critical therapeutic intervention in CRC patients. The demonstrated power of this research strategy makes it generally applicable to other physiological and pathological situations.

#### Usage

```
data( GSE3964_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Graudens E, Boulanger V, Mollard C, Mariage-Samson R et al.??Deciphering cellular s
 Laboratory: Graudens, Imbeaud 2006
 Contact information:
 Title: Deciphering cellular states of innate tumor drug responses.
 URL:
 PMIDs: 16542501
  Abstract: A 242 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      11K_VJF-ARRAY
   platform_shorttitle:
   platform_summary:
      11k_vjf-array
   platform_manufacturer:
      Array s/IMAGE - Genexpress
   platform_distribution:
      non-commercial
   platform_accession:
      GPL3282
   platform_technology:
      spotted DNA/cDNA
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 384D8-2 38600 ... ZYX (5845 total)
```

GSE3964\_eset 77

varLabels: probeset gene
varMetadata: labelDescription

#### **Details**

```
assayData: 5845 features, 29 samples
Platform type: 11k_vjf-array
_____
Available sample meta-data:
alt_sample_name:
  Length
          Class
                      Mode
      29 character character
sample_type:
adjacentnormal
              metastatic
                                   tumor
           6
                  14
primarysite:
СО
29
summarystage:
late
 29
M:
1
29
Dstage:
D
29
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median
                      Mean 3rd Qu.
                                       Max.
 48.00 60.00 62.00
                        61.31 68.00
                                     71.00
summarylocation:
1
29
gender:
f m
11 18
stageall:
```

78 *GSE4045\_eset* 

```
4
29
drug_name:
FOLFIRI
     29
drug_treatment:
29
drug_response:
n y
15 14
preop_drug_treatment:
 n
29
fu:
У
29
irinotecan:
У
29
folfiri:
У
29
leucovorin:
 У
29
chemotherapy:
29
uncurated\_author\_metadata:
   Length
             Class
       29 character character
```

GSE4045\_eset

Serrated carcinomas form a subclass of colorectal cancer with distinct molecular basis.

*GSE4045\_eset* 79

#### **Description**

Serrated colorectal carcinomas (CRCs) are morphologically different from conventional CRCs and have been proposed to follow a distinct pathway of CRC formation. Despite studies of single molecular events in this tumor type, the diagnosis of serrated CRC relies on morphology and the putative unique biological character of these tumors has not been established. Here we show that the gene expression profiling of 37 CRCs separated serrated and conventional CRCs into two distinct branches in unsupervised hierarchical clustering (P-value 7.8 x 10(-7)), and revealed 201 differentially expressed genes representing potential biomarkers for serrated CRC. Immunohistochemistry was utilized to verify the key findings in the 37 CRCs examined by expression profiling, and a separate validation set of 37 serrated and 86 conventional CRCs was examined to evaluate the candidate biomarkers in an extended sample material. Ephrin receptor B2, hypoxia-inducible factor 1-alpha and patched appeared as proteins important for genesis of serrated CRC. This study establishes serrated CRCs as a biologically distinct subclass of CRC and represents a step forward in the molecular classification of these cancers. The study also provides a platform to understand the molecular basis of serrated CRC and in long term may contribute to the development of specific treatment options for this tumor type.

#### Usage

```
data( GSE4045_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Laiho P, Kokko A, Vanharanta S, Salovaara R et al.??Serrated carcinomas form a subcl
 Laboratory: Laiho, Aaltonen 2007
 Contact information:
 Title: Serrated carcinomas form a subclass of colorectal cancer with distinct molecular basis.
 URL:
 PMIDs: 16819509
 Abstract: A 205 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133A] Affymetrix Human Genome U133A Array
  platform_shorttitle:
     Affymetrix HG-U133A
  platform_summary:
     hgu133a
  platform_manufacturer:
     Affymetrix
  platform_distribution:
     commercial
  platform_accession:
     GPL96
  platform_technology:
```

80  $GSE4045\_eset$ 

```
in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: frma
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: A1CF A2M ... ZZZ3 (12986 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 12986 features, 37 samples
   Platform type: hgu133a
   _____
```

```
Available sample meta-data:
_____
alt_sample_name:
  Length
         Class
                     Mode
      37 character character
sample_type:
tumor
  37
summarygrade:
low NA's
 36
G:
  1
      2 3 NA's
  4
     28
           4 1
summarystage:
early late
   2
       35
M:
0 1
35 2
Dstage:
B C D
2 33 2
family_history:
```

GSE4526\_eset 81

```
y NA's
   n
  32
msi:
MSI MSS
  1 36
location:
  distal proximal
      22
               15
summarylocation:
1 r
22 15
gender:
f m
19 18
stageall:
   3
        4 NA's
  33
        2
             2
batch:
                          Mode
              Class
   Length
       37 character character
uncurated_author_metadata:
   Length
              Class
                          Mode
       37 character character
```

GSE4526\_eset

Gene expression signature for recurrence in stage III colorectal cancers.

#### **Description**

Colorectal cancer patients with lymph node metastases (stage III) show poorer prognosis than those without. Predicting development of recurrence may guide the need for intensive follow-up and/or adjuvant chemotherapy in such patients. The authors' objective was to identify a set of discriminating genes that could predict recurrence in stage III colorectal cancer. Thirty-six stage III colorectal cancer patients were studied. Tumor samples were obtained from surgically resected specimens. Thirteen patients developed recurrence, whereas 23 patients did not. Gene expression profiles were determined using human HG-U133 Plus 2.0 Gene Chip (Affymetrix, Santa Clara, Calif). The authors identified 45 discriminating genes between patients with and without recurrence. By using this gene set, they established a new model to predict recurrence with an accuracy of 90.9%. The

82 *GSE4526\_eset* 

discriminating genes included calcineurin-binding protein 1 (CABIN1), whose expression differed remarkably between patients with and without recurrence (P=.0073). The authors further examined the DNA copy number of CABIN1 and were able to show a significant relation with recurrence (P<.012). Patients having CABIN1 gene loss demonstrated a higher risk of recurrence (odds ratio, 18.8). DNA copy number of CABIN1 alone could predict recurrence with an accuracy of 80.0%. The results of the current study demonstrated that gene expression profiling is useful in predicting recurrence in stage III colorectal cancer. The authors identified CABIN1 among discriminating genes that may play a key role in the development of recurrence. These results may help to establish an individualized therapy for stage III colorectal cancer. Copyright (c) 2009 American Cancer Society.

#### Usage

```
data( GSE4526_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Watanabe T, Kobunai T, Sakamoto E, Yamamoto Y et al.??Gene expression signature for
 Laboratory: Watanabe T,??Kobunai T,??Toda E,??Oka T 2006
 Contact information:
 Title: Gene expression signature for recurrence in stage III colorectal cancers.
 URL:
 PMIDs: 19016304
  Abstract: A 246 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
```

GSE4526\_eset 83

varLabels: probeset gene
varMetadata: labelDescription

#### **Details**

```
assayData: 19320 features, 36 samples
Platform type: hgu133plus2
_____
Available sample meta-data:
alt_sample_name:
         Class Mode
  Length
      36 character character
sample_type:
tumor
  36
summarystage:
late
 36
M:
0
36
Dstage:
С
36
recurrence_status:
norecurrence recurrence
        23 13
stageall:
3
36
ethnicity:
other
  36
preop_drug_treatment:
36
uncurated_author_metadata:
```

84 *GSE45270\_eset* 

```
Length Class Mode
36 character character
```

GSE45270\_eset

AMC tubular and serrated adenomas

### Description

Profiling project of a panel of tubular adenoma and serrated adenoma patient material collected in the Academic Medical Center

#### Usage

```
data( GSE45270_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Profiling project of a panel of tubular adenoma and serrated adenoma patient materia
 Laboratory: Medema JP,??de Sousa E Melo F,??Vermeulen L,??Jansen M,??Dekker E,??Van Noesel C,??Fesslen
 Contact information:
 Title: AMC tubular and serrated adenomas
 URL:
 PMIDs: PMID unknown
 Abstract: A 19 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      [HG-U133_Plus_2] Affymetrix Human Genome U133 Plus 2.0 Array
   platform_shorttitle:
      Affymetrix HG-U133Plus2
   platform_summary:
      hgu133plus2
   platform_manufacturer:
      Affymetrix
   platform_distribution:
      commercial
   platform_accession:
      GPL570
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
```

```
Preprocessing: frma
featureData(eset):
An object of class 'AnnotatedDataFrame'
featureNames: A1BG A1BG-AS1 ... ZZZ3 (19320 total)
varLabels: probeset gene
varMetadata: labelDescription
```

#### **Details**

```
assayData: 19320 features, 13 samples
Platform type: hgu133plus2
Available sample meta-data:
 -----
alt_sample_name:
  Length
             Class
                       Mode
      13 character character
batch:
2012-08-23
       13
uncurated_author_metadata:
  Length
             Class
                       Mode
      13 character character
```

TCGA.COAD\_eset Comprehensive molecular characterization of human colon and rectal

cancer.

#### **Description**

To characterize somatic alterations in colorectal carcinoma, we conducted a genome-scale analysis of 276 samples, analysing exome sequence, DNA copy number, promoter methylation and messenger RNA and microRNA expression. A subset of these samples (97) underwent low-depth-of-coverage whole-genome sequencing. In total, 16% of colorectal carcinomas were found to be hypermutated: three-quarters of these had the expected high microsatellite instability, usually with hypermethylation and MLH1 silencing, and one-quarter had somatic mismatch-repair gene and polymerase?? (POLE) mutations. Excluding the hypermutated cancers, colon and rectum cancers were found to have considerably similar patterns of genomic alteration. Twenty-four genes were significantly mutated, and in addition to the expected APC, TP53, SMAD4, PIK3CA and KRAS mutations, we found frequent mutations in ARID1A, SOX9 and FAM123B. Recurrent copy-number alterations include potentially drug-targetable amplifications of ERBB2 and newly discovered amplification of IGF2. Recurrent chromosomal translocations include the fusion of NAV2 and WNT

pathway member TCF7L1. Integrative analyses suggest new markers for aggressive colorectal carcinoma and an important role for MYC-directed transcriptional activation and repression.

#### Usage

```
data( TCGA.COAD_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Comprehensive molecular characterization of human colon and rectal cancer. Nature 2
 Laboratory: The Cancer Genome Atlas Network 2012
 Contact information:
 Title: Comprehensive molecular characterization of human colon and rectal cancer.
 URI:
 PMIDs: 22810696
 Abstract: A 168 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      Agilent 244K Custom Gene Expression G4502A-07-3
   platform_shorttitle:
      Agilent G4502A-07-3
   platform_summary:
      agilent-014850 whole human genome microarray 4x44k g4112f
   platform_manufacturer:
      Agilent
   platform_distribution:
      commercial
   platform_accession:
      NA
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 15E1.2 2'-PDE ... ZZZ3 (17814 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

#### **Details**

assayData: 17814 features, 130 samples

```
Platform type: agilent-014850 whole human genome microarray 4x44k g4112f
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)
  124 observations deleted due to missingness
records
        n.max n.start events median 0.95LCL 0.95UCL
  6.00
         6.00 6.00
                      5.00
                             1.37 1.16
-----
Available sample meta-data:
-----
unique_patient_ID:
           Class
  Length
                      Mode
     130 character character
alt_sample_name:
  Length
                      Mode
     130 character character
sample_type:
adjacentnormal
                     tumor
           7
                       123
primarysite:
 co NA's
129
     1
summarystage:
early late NA's
  75
       54
             1
T:
1 2 3 4
3 27 88 12
N:
  0
      1
          2 NA's
 80
      22
         27 1
M:
       1 NA's
  0
108
      21 1
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median Mean 3rd Qu.
                                      Max.
  36.00 65.00 72.00
                      70.89 80.00
                                      90.00
```

```
Min. 1st Qu. Median Mean 3rd Qu. Max. NA's
  29.0 442.0 500.0 494.5 681.0 774.0 124
vital_status:
deceased living NA's
 5 15 110
msi:
MSI MSS NA's
 1 4 125
location:
 Length Class Mode
   130 character character
summarylocation:
 l r NA's
 59 70 1
gender:
f m
67 63
kras:
mutant wt NA's 2 1 127
stageall:
 1 2 3 4 NA's
 25 51 30 20 4
ethnicity:
   black caucasian NA's 3 17 110
drug_treatment:
 n y NA's
  2 3 125
preop_drug_treatment:
n y
127 3
fu:
 n NA's
 31 99
```

days\_to\_death:

## bevacizumab: n NA's 31 99 irinotecan: n NA's 31 99 cetuximab: n NA's 31 99 capecitabine: n NA's 31 99 dexamethasone: n NA's 31 99 gcsf: n NA's 31 99 fudr: n NA's 31 99 folfiri: n NA's 31 99 folfox: n NA's 31 99 leucovorin: n NA's 31 99 mitomycin: n y NA's 30 1 99

platin: n NA's 31 99

```
panitumumab:
   n NA's
  31
       99
pegfilgrastim:
   n NA's
  31
       99
raltitrexed:
   n NA's
  31
       99
ancillary:
        y NA's
  30
        1
             99
chemotherapy:
   y NA's
  31
       99
moltherapy:
        y NA's
   n
             99
  21
       10
uncurated_author_metadata:
   Length
               Class
                           Mode
      130 character character
```

TCGA.READ\_eset

Comprehensive molecular characterization of human colon and rectal cancer.

### Description

To characterize somatic alterations in colorectal carcinoma, we conducted a genome-scale analysis of 276 samples, analysing exome sequence, DNA copy number, promoter methylation and messenger RNA and microRNA expression. A subset of these samples (97) underwent low-depth-of-coverage whole-genome sequencing. In total, 16% of colorectal carcinomas were found to be hypermutated: three-quarters of these had the expected high microsatellite instability, usually with hypermethylation and MLH1 silencing, and one-quarter had somatic mismatch-repair gene and polymerase?? (POLE) mutations. Excluding the hypermutated cancers, colon and rectum cancers were found to have considerably similar patterns of genomic alteration. Twenty-four genes were significantly mutated, and in addition to the expected APC, TP53, SMAD4, PIK3CA and KRAS mutations, we found frequent mutations in ARID1A, SOX9 and FAM123B. Recurrent copy-number alterations include potentially drug-targetable amplifications of ERBB2 and newly discovered amplification of IGF2. Recurrent chromosomal translocations include the fusion of NAV2 and WNT

pathway member TCF7L1. Integrative analyses suggest new markers for aggressive colorectal carcinoma and an important role for MYC-directed transcriptional activation and repression.

#### Usage

```
data( TCGA.READ_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Comprehensive molecular characterization of human colon and rectal cancer. Nature 2
 Laboratory: The Cancer Genome Atlas Network 2012
 Contact information:
 Title: Comprehensive molecular characterization of human colon and rectal cancer.
 URI:
 PMIDs: 22810696
 Abstract: A 168 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
  platform_title:
      Agilent 244K Custom Gene Expression G4502A-07-3
   platform_shorttitle:
      Agilent G4502A-07-3
   platform_summary:
      agilent-014850 whole human genome microarray 4x44k g4112f
   platform_manufacturer:
      Agilent
   platform_distribution:
      commercial
   platform_accession:
      NA
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: 15E1.2 2'-PDE ... ZZZ3 (17814 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

#### Details

assayData: 17814 features, 51 samples

```
Platform type: agilent-014850 whole human genome microarray 4x44k g4112f
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)
  50 observations deleted due to missingness
records
         n.max n.start events median 0.95LCL 0.95UCL
 1.000
        1.000 1.000 1.000 0.866
-----
Available sample meta-data:
-----
unique_patient_ID:
           Class
  Length
                      Mode
      51 character character
alt_sample_name:
  Length
                      Mode
      51 character character
sample_type:
tumor
  51
primarysite:
 re NA's
 50 1
summarystage:
early late
  30
       21
T:
1 2 3 4
3 9 37 2
N:
0 1 2
30 11 10
M:
0 1
age_at_initial_pathologic_diagnosis:
  Min. 1st Qu. Median Mean 3rd Qu.
                                       Max.
 41.00 62.00 67.00 67.67 73.50
                                      89.00
```

```
days_to_tumor_recurrence:
301 327 NA's
 1 1 49
recurrence_status:
                         NA's
norecurrence recurrence
                           46
            2
days_to_death:
316 NA's
 1 50
vital_status:
deceased living NA's
    1 4
                 46
msi:
MSS NA's
 1 50
location:
rectosigmoid
             rectum
                         NA's
                           1
  3
                47
summarylocation:
 l NA's
 50 1
gender:
f m
24 27
kras:
mutant
        wt NA's
    2
         2
            47
lymphnodesremoved:
  Min. 1st Qu. Median Mean 3rd Qu.
                                 Max.
 12.00 17.00 22.00
                    24.73 32.00 65.00
lymphnodesinvaded:
  Min. 1st Qu. Median
                    Mean 3rd Qu.
                                Max.
 0.000 0.000 0.000 2.608 3.000 31.000
stageall:
 1 2
        3 4 NA's
 12 18 13 7 1
```

```
black caucasian NA's
      1 4
                      46
drug_treatment:
  y NA's
  2 49
preop_drug_treatment:
51
fu:
 n NA's
 16 35
bevacizumab:
 n NA's
 16 35
irinotecan:
  n NA's
 16 35
cetuximab:
  n NA's
 16 35
capecitabine:
 n NA's
 16 35
dexamethasone:
  n NA's
 16 35
gcsf:
 n NA's
 16 35
fudr:
  n NA's
 16 35
folfiri:
 n NA's
```

16 35

ethnicity:

folfox:

```
n NA's
 16 35
leucovorin:
  n NA's
 16 35
mitomycin:
  n NA's
 16 35
platin:
  n NA's
 16 35
panitumumab:
  n NA's
 16 35
pegfilgrastim:
  n NA's
 16 35
raltitrexed:
  n NA's
 16 35
ancillary:
  n NA's
 17 34
chemotherapy:
  y NA's
 17 34
moltherapy:
  n
     y NA's
 15
       2 34
uncurated_author_metadata:
  Length
            Class
                       Mode
      51 character character
```

TCGA.RNASeqV2.READ\_eset

Comprehensive molecular characterization of human colon and rectal cancer.

#### **Description**

To characterize somatic alterations in colorectal carcinoma, we conducted a genome-scale analysis of 276 samples, analysing exome sequence, DNA copy number, promoter methylation and messenger RNA and microRNA expression. A subset of these samples (97) underwent low-depth-of-coverage whole-genome sequencing. In total, 16% of colorectal carcinomas were found to be hypermutated: three-quarters of these had the expected high microsatellite instability, usually with hypermethylation and MLH1 silencing, and one-quarter had somatic mismatch-repair gene and polymerase?? (POLE) mutations. Excluding the hypermutated cancers, colon and rectum cancers were found to have considerably similar patterns of genomic alteration. Twenty-four genes were significantly mutated, and in addition to the expected APC, TP53, SMAD4, PIK3CA and KRAS mutations, we found frequent mutations in ARID1A, SOX9 and FAM123B. Recurrent copy-number alterations include potentially drug-targetable amplifications of ERBB2 and newly discovered amplification of IGF2. Recurrent chromosomal translocations include the fusion of NAV2 and WNT pathway member TCF7L1. Integrative analyses suggest new markers for aggressive colorectal carcinoma and an important role for MYC-directed transcriptional activation and repression.

#### Usage

```
data( TCGA.RNASeqV2.READ_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Comprehensive molecular characterization of human colon and rectal cancer. Nature 2
 Laboratory: The Cancer Genome Atlas Network 2012
 Contact information:
 Title: Comprehensive molecular characterization of human colon and rectal cancer.
 URL:
 PMIDs: 22810696
  Abstract: A 168 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [RNASeqV2] Illumina HiSeq RNA sequencing
   platform_shorttitle:
   platform_summary:
   platform_manufacturer:
      Illumina
   platform_distribution:
```

```
sequencing
      platform_accession:
      platform_technology:
         in situ oligonucleotide
      warnings:
         No warnings yet
   Preprocessing: default
   featureData(eset):
   An object of class 'AnnotatedDataFrame'
     featureNames: ? A1BG ... ZZZ3 (20502 total)
     varLabels: probeset gene
     varMetadata: labelDescription
Details
   assayData: 20502 features, 6 samples
   Platform type: NA
   Overall survival time-to-event summary (in years):
   Call: survfit(formula = Surv(time, cens) ~ -1)
      3 observations deleted due to missingness
   records n.max n.start events median 0.95LCL 0.95UCL
      3.00
            3.00 3.00 3.00 3.44
                                             2.72
   Available sample meta-data:
   unique_patient_ID:
      Length Class
                            Mode
           6 character character
   alt_sample_name:
      Length
                Class
                            Mode
           6 character character
   sample_type:
   tumor
       6
   primarysite:
   re
    6
   summarystage:
   early late
```

```
3 3
T:
2 3
1 5
N:
0 1 2
3 2 1
M:
0
{\tt age\_at\_initial\_pathologic\_diagnosis:}
56 57 72 73 77
1 1 1 1 2
days_to_tumor_recurrence:
630 3316 NA's
1 1 4
recurrence_status:
days_to_death:
991 1257 1741 NA's
 1 1 1 3
vital_status:
deceased living NA's
  3 2 1
msi:
MSS NA's
5 1
location:
rectosigmoid rectum
summarylocation:
1
6
gender:
```

```
6
kras:
mutant
           wt
     3
            3
lymphnodesremoved:
7 14 15 16
1 2 1 2
lymphnodesinvaded:
0 1 5
3 2 1
stageall:
2 3
3 3
ethnicity:
caucasian
        6
preop_drug_treatment:
n
6
uncurated_author_metadata:
   Length
              Class
                          Mode
        6 character character
```

TCGA.RNASeqV2\_eset

Comprehensive molecular characterization of human colon and rectal cancer.

#### Description

To characterize somatic alterations in colorectal carcinoma, we conducted a genome-scale analysis of 276 samples, analysing exome sequence, DNA copy number, promoter methylation and messenger RNA and microRNA expression. A subset of these samples (97) underwent low-depth-of-coverage whole-genome sequencing. In total, 16% of colorectal carcinomas were found to be hypermutated: three-quarters of these had the expected high microsatellite instability, usually with hypermethylation and MLH1 silencing, and one-quarter had somatic mismatch-repair gene and polymerase ?? (POLE) mutations. Excluding the hypermutated cancers, colon and rectum cancers were found to have considerably similar patterns of genomic alteration. Twenty-four genes were significantly mutated, and in addition to the expected APC, TP53, SMAD4, PIK3CA and KRAS mutations, we found frequent mutations in ARID1A, SOX9 and FAM123B. Recurrent copy-number

alterations include potentially drug-targetable amplifications of ERBB2 and newly discovered amplification of IGF2. Recurrent chromosomal translocations include the fusion of NAV2 and WNT pathway member TCF7L1. Integrative analyses suggest new markers for aggressive colorectal carcinoma and an important role for MYC-directed transcriptional activation and repression.

#### Usage

```
data( TCGA.RNASeqV2_eset )
```

#### **Format**

```
experimentData(eset):
Experiment data
 Experimenter name: Comprehensive molecular characterization of human colon and rectal cancer. Nature 2
  Laboratory: The Cancer Genome Atlas Network 2012
  Contact information:
 Title: Comprehensive molecular characterization of human colon and rectal cancer.
  URL:
  PMIDs: 22810696
  Abstract: A 168 word abstract is available. Use 'abstract' method.
  Information is available on: preprocessing
  notes:
   platform_title:
      [RNASeqV2] Illumina HiSeq RNA sequencing
   platform_shorttitle:
   platform_summary:
   platform_manufacturer:
      Illumina
   platform_distribution:
      sequencing
   platform_accession:
   platform_technology:
      in situ oligonucleotide
   warnings:
      No warnings yet
Preprocessing: default
featureData(eset):
An object of class 'AnnotatedDataFrame'
  featureNames: ? A1BG ... ZZZ3 (20502 total)
  varLabels: probeset gene
  varMetadata: labelDescription
```

#### **Details**

```
assayData: 20502 features, 195 samples
Platform type: NA
Overall survival time-to-event summary (in years):
Call: survfit(formula = Surv(time, cens) ~ -1)
  174 observations deleted due to missingness
records n.max n.start events median 0.95LCL 0.95UCL
21.000 21.000 21.000 18.000 1.208 0.715 5.605
Available sample meta-data:
_____
unique_patient_ID:
  Length
          Class
                       Mode
     195 character character
alt_sample_name:
  Length
           Class
                      Mode
     195 character character
sample_type:
                    tumor
adjacentnormal
          14
                      181
primarysite:
СО
195
summarystage:
early late NA's
  91 103 1
T:
  Length
            Class
                       Mode
     195 character character
N:
 0
    1 2 X
116 51 27 1
M:
  0
      1
           X NA's
128
      29
           35 3
{\tt age\_at\_initial\_pathologic\_diagnosis:}
  Min. 1st Qu. Median Mean 3rd Qu.
                                        Max.
```

31.00 55.00 66.00 64.95 77.00 90.00

days\_to\_tumor\_recurrence:

Min. 1st Qu. Median Mean 3rd Qu. Max. NA's 0.0 306.2 458.5 539.7 601.5 2564.0 169

recurrence\_status:

norecurrence recurrence NA's 111 25 59

days\_to\_death:

Min. 1st Qu. Median Mean 3rd Qu. Max. NA's 43.0 187.0 424.0 816.6 1422.0 2134.0 174

vital\_status:

deceased living NA's 18 119 58

msi:

MSI MSS NA's 9 32 154

location:

Length Class Mode 195 character character

summarylocation:

1 r NA's 65 127 3

gender:

f m 87 108

kras:

mutant wt NA's 4 9 182

lymphnodesremoved:

Min. 1st Qu. Median Mean 3rd Qu. Max. NA's 0.00 13.25 19.00 23.09 27.00 109.00 21

lymphnodesinvaded:

Min. 1st Qu. Median Mean 3rd Qu. Max. NA's 0.000 0.000 0.000 1.983 2.000 50.000 22

stageall:

1 2 3 4 NA's

folfiri:

n y NA's

```
33 73 52 30 7
ethnicity:
                   other NA's
12 15
   black caucasian
     16 152
drug_treatment:
  n y NA's
 21 14 160
{\tt preop\_drug\_treatment:}
195
fu:
 n NA's
 60 135
bevacizumab:
  n NA's
 60 135
irinotecan:
  n NA's
 60 135
cetuximab:
  n NA's
 60 135
capecitabine:
  n NA's
 60 135
dexamethasone:
  n NA's
 60 135
gcsf:
  n NA's
 60 135
fudr:
 n y NA's
 57 3 135
```

59 1 135

folfox:

n NA's

60 135

leucovorin:

n NA's

60 135

mitomycin:

n NA's

60 135

platin:

n y NA's

58 2 135

panitumumab:

n NA's

60 135

pegfilgrastim:

n y NA's

59 1 135

raltitrexed:

n y NA's

59 1 135

ancillary:

n y NA's

58 2 135

chemotherapy:

n y NA's

2 58 135

moltherapy:

n y NA's

57 3 135

 $uncurated\_author\_metadata:$ 

Length Class Mode

195 character character

# **Index**

| * datasets                       | GSE13067_eset (GSE13067_eset), 12   |
|----------------------------------|-------------------------------------|
| GSE11237_eset, 3                 | GSE13294_eset (GSE13294_eset), 14   |
| GSE12225.GPL3676_eset, 6         | GSE14095_eset (GSE14095_eset), 16   |
| GSE12945_eset, 9                 | GSE14333_eset (GSE14333_eset), 18   |
| GSE13067_eset, 12                | GSE16125.GPL5175_eset               |
| GSE13294_eset, 14                | (GSE16125.GPL5175_eset), 21         |
| GSE14095_eset, 16                | GSE17536_eset (GSE17536_eset), 25   |
| GSE14333_eset, 18                | GSE17537_eset (GSE17537_eset), 28   |
| GSE16125.GPL5175_eset, 21        | GSE17538.GPL570_eset                |
| GSE17536_eset, 25                | (GSE17538.GPL570_eset), 32          |
| GSE17537_eset, 28                | GSE18105_eset (GSE18105_eset), 35   |
| GSE17538.GPL570_eset, 32         | GSE2109_eset (GSE2109_eset), 38     |
| GSE18105_eset, 35                | GSE21510_eset (GSE21510_eset), 40   |
| GSE2109_eset, 38                 | GSE21815_eset (GSE21815_eset), 43   |
| GSE21510_eset, 40                | GSE24549.GPL5175_eset               |
| GSE21815_eset, 43                | (GSE24549.GPL5175_eset), 45         |
| GSE24549.GPL5175_eset, 45        | GSE24550.GPL5175_eset               |
| GSE24550.GPL5175_eset,48         | (GSE24550.GPL5175_eset), 48         |
| GSE2630_eset, 51                 | GSE2630_eset (GSE2630_eset), 51     |
| GSE26682.GPL570_eset, 54         | GSE26682.GPL570_eset                |
| GSE26682.GPL96_eset, 56          | (GSE26682.GPL570_eset), 54          |
| GSE26906_eset, 59                | GSE26682.GPL96_eset                 |
| GSE27544_eset, 62                | (GSE26682.GPL96_eset), 56           |
| GSE28702_eset, 63                | GSE26906_eset (GSE26906_eset), 59   |
| GSE3294_eset, 66                 | GSE27544_eset (GSE27544_eset), 62   |
| GSE33113_eset, 69                | GSE28702_eset (GSE28702_eset), 63   |
| GSE39582_eset, 72                | GSE3294_eset (GSE3294_eset), 66     |
| GSE3964_eset, 75                 | GSE33113_eset (GSE33113_eset), 69   |
| GSE4045_eset, 78                 | GSE39582_eset (GSE39582_eset), 72   |
| GSE4526_eset, 81                 | GSE3964_eset (GSE3964_eset), 75     |
| GSE45270_eset, 84                | GSE4045_eset (GSE4045_eset), 78     |
| TCGA.COAD_eset, 85               | GSE4526_eset (GSE4526_eset), 81     |
| TCGA.READ_eset, 90               | GSE45270_eset (GSE45270_eset), 84   |
| TCGA.RNASeqV2.READ_eset, 96      | TCGA.COAD_eset (TCGA.COAD_eset), 85 |
| TCGA.RNASeqV2_eset, 99           | TCGA.READ_eset (TCGA.READ_eset), 90 |
| GSE11237_eset (GSE11237_eset), 3 | TCGA.RNASeqV2.READ_eset             |
| GSE12225.GPL3676_eset            | (TCGA.RNASeqV2.READ_eset), 96       |
| (GSE12225.GPL3676_eset), 6       | TCGA.RNASeqV2_eset                  |
| GSE12945_eset (GSE12945_eset), 9 | (TCGA.RNASeqV2_eset), 99            |
| 00212070_0000 (00212070_0000), / | (100/1.111/Jeq12_e3et), 99          |

106 INDEX

```
curatedCRCData
        (curatedCRCData-package), 2
curatedCRCData-package, 2
GSE11237_eset, 3
GSE12225.GPL3676_eset, 6
GSE12945_eset, 9
GSE13067_eset, 12
GSE13294_eset, 14
GSE14095_eset, 16
GSE14333_eset, 18
GSE16125.GPL5175_eset, 21
GSE17536_eset, 25
GSE17537_eset, 28
GSE17538.GPL570_eset, 32
GSE18105_eset, 35
GSE2109_eset, 38
GSE21510_eset, 40
GSE21815_eset, 42
GSE24549.GPL5175_eset, 45
GSE24550.GPL5175_eset, 48
GSE2630_eset, 51
GSE26682.GPL570_eset, 54
GSE26682.GPL96_eset, 56
GSE26906_eset, 59
GSE27544_eset, 62
GSE28702_eset, 63
GSE3294_eset, 66
GSE33113_eset, 69
GSE39582_eset, 72
GSE3964_eset, 75
GSE4045_eset, 78
GSE4526_eset, 81
GSE45270_eset, 84
TCGA.COAD_eset, 85
TCGA.READ_eset, 90
TCGA.RNASeqV2.READ_eset, 95
TCGA.RNASeqV2_eset, 99
```